Document Identifier: **PMC8476158**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
Six samples were collected in the native Asian area , more precisely in four Chinese and two Japanese localities .
Sex as a biological variable
not required.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
To test whether the candidate regions were enriched in TEs, we generated random expectations by applying the above to 1,000 randomly selected SNPs 250 times.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The previously established consensus were obtained by extracting all Drosophila consensus from Dfam and Repbase databases (release 2016-2018 for both) (Hubley et al. 2016 https:www.girinst.orgrepbase, last accessed December 01, 2019).	Repbase		Suggestion: Repbase (RRID:SCR_021169)(link)
Graphical visualization of the results was produced using Circos (Krzywinski et al. 2009) .	Circos		Suggestion: Circos (RRID:SCR_011798)(link)
Evaluation of Population Genetics Statistics We estimated Wattersons theta (th w) and Tajimas D statistics in nonoverlapping 1,000 bp windows using PoPoolation (v1.2.2) (Kofler et al. 2011) .	PoPoolation		Suggestion: PoPoolation (RRID:SCR_003495)(link)
Reads were filtered for a minimum mapping quality of 20 and a pileup file generated with samtools (v1.7) (Li et al. 2009) .	samtools		
For each region , we looked for overlapping TEs using the RepeatMasker annotation (gff file , see Annotation of the reference genome) .	RepeatMasker		Suggestion: RepeatMasker (RRID:SCR_012954)(link)
Second , we retrieved homologous regions in the D . melanogaster genome using BLAT against the D . melanogaster masked assembly downloaded from UCSC Genome Browser (http:hgdownload.soe.ucsc.edugoldenPathdm6bigZips , last accessed December 01	BLAT		Suggestion: BLAT (RRID:SCR_011919)(link)
	UCSC Genome Browser		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Correlation between Climatic Variables and TE Family Abundance Partial Mantel tests were used to test the correlation between bioclimatic variables and TE family abundance correcting for population structure (as in Quadrana et al. 2016).	Mantel tests
P values were further adjusted to account for multiple testing applying the Benjamini-Hochberg correction (Benjamini and Hochberg 1995).	Benjamini-Hochberg
	Benjamini and Hochberg

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq