Document Identifier: **PMC8480666**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: All the experiments were approved by the local ethics committee (APAFIS , reference 2018070316423884) , and experiments were conducted by authorized personnel in the in-house specific pathogen–free animal facility (agreement reference B75-13-19; CEEA-005) .
Consent: All patients included in the present study gave their consent for DNA testing in the research setting , and consent forms (RBM 01-29) were signed by each participant .
Euthanasia Agents: Briefly , mice were killed by CO2 asphyxiation and enucleated.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: Lipidomic analyses (Colsch et al. , 2015) 8-mo-old Cyp2u1+/+ , Cyp2u1 +/– , and Cyp2u1 – / – mice were killed by CO2 asphyxiation , and the cerebral cortices were immediately dissected and frozen in isopentane.
Randomization
Bias was minimized by randomly testing mice in experimental cohorts , regardless of their genotype.
Blinding
not detected.
Power Analysis
not detected.

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
For immunohistochemical analyses , after 90-min incubation in blocking solution , sections were incubated with primary antibodies (mouse anti-NeuN; MAB377; Merck Millipore) , rabbit anti-glial fibrillary acidic protein (Z0334; Dako) , and mouse anti-calbindin (300; Swant) in 2 % BSA/0.25 % Triton X-100 in PBS overnight at 4°C .	anti-NeuN		
	anti-glial fibrillary acidic protein (Z0334; Dako) ,		
	anti-calbindin		
Retinal and RPE/choroid tissues were dissected intact from the globe , flat-mounted , and processed for immunohistochemistry using polyclonal goat anti-IBA1 (1:500 , ab5076; Merck Millipore) , rabbit anti-opsin red/green (1:200 , AB5405; Merck Millipore) , rabbit anti-opsin blue (1:200 , ab5407; Merck Millipore) , rabbit anti-cone arrestin (ab15282; Merck Millipore) , and donkey anti-rabbit secondary antibody conjugated with Alexa Fluor 594 and donkey anti-goat secondary antibody conjugated with Alexa Fluor 488 (Life Technologies) .	anti-IBA1		
	anti-opsin red/ green		
	anti-opsin blue		
	anti-cone arrestin		
	anti-rabbit		
	anti-goat		
The following primary antibodies were used: rabbit anti-α-actin (1:5,000 , ab15246; Abcam) , mouse anti-COQ6 (1:100 , sc-393932; Santa Cruz Biotechnology) , mouse anti-COQ9 (1:1,000 , sc-271892; Santa Cruz Biotechnology) , mouse anti-COQ5 (1:500 , 17453-1-AP; Proteintech) , and mouse anti-OXPHOS antibody cocktail (1:1,000 , ab110413; Abcam) .	anti-α-actin		
	anti-COQ6		
	anti-COQ9		
	anti-COQ5		
	anti-OXPHOS		
The following secondary antibodies were used: donkey anti-rabbit IgG (NA9340V; GE Healthcare) and sheep anti-mouse (NXA931; GE Healthcare) .	anti-rabbit IgG		
	anti-mouse		
Experimental Models: Organisms/Strains			
Materials and methods Mice All mice were maintained on a C57BL/6N genetic background.	C57BL/6N		Suggestion: RRID:IMSR_RJ:C57BL-6NRJ(link)
The concentration was calculated to deliver 7,500 mg sodium formate/kg/ d , based on an average water intake	C57BL/6		Suggestion: RRID:IMSR_JAX:000664(link)

of 5 ml/d for a 25-g C57BL/6 mouse (Bryant et al. , 2018; Momb et al. , 2013) .			
Lipidomic analyses (Colsch et al. , 2015) 8-mo-old Cyp2u1+/+ , Cyp2u1 +/– , and Cyp2u1 – / – mice were killed by CO2 asphyxiation , and the cerebral cortices were immediately dissected and frozen in isopentane.	Cyp2u1 – / –		
Software and Algorithms			
Gait was analyzed using GaitScan software (CleverSys) .	GaitScan		
Counting of cell populations was performed using ImageJ software.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)

Other Entities Detected

Your Sentences	Recognized Entity
Oligonucleotides	
It was performed by PCR with the 5#-GTAGGAGAGGTCTCTACAGAGGTG-3# and 5#-CCAACAGCTTCCCCACAACGG-3# primers to amplify a 292-bp amplicon from the KI allele.	5#-GTAGGAGAGGTCTCTACAGAGGTG-3#
	5#-CCAACAGCTTCCCCACAACGG-3#
To detect the wild-type allele , the primers 5#-GGGTAAAGGCAGGTTCTTTGAACCC-3# and 5#-CCTCTCTACACGCCACTTCTCTAAC-3# were used.	5#-GGGTAAAGGCAGGTTCTTTGAACCC-3#
	5#-CCTCTCTACACGCCACTTCTCTAAC-3#
Cre recombinase was detected using 5#-GCATTACCGGTCGATGCAACGAGTGATGAG-3# and 5#-GAGTGAACGAACCTGGTCGAAATCAGTGCG-3# to produce a 408-bp amplicon .	5#-GCATTACCGGTCGATGCAACGAGTGATGAG-3#
	5#-GAGTGAACGAACCTGGTCGAAATCAGTGCG-3#
Unresolved RRIDs	
Retinal and RPE/choroid tissues were dissected intact from the globe , flat-mounted , and processed for immunohistochemistry using polyclonal goat anti-IBA1 (1:500 , ab5076; Merck Millipore) , rabbit anti-opsin red/green (1:200 , AB5405; Merck Millipore) , rabbit anti-opsin blue (1:200 , ab5407; Merck Millipore) , rabbit anti-cone arrestin (ab15282; Merck Millipore) , and donkey anti-rabbit secondary antibody conjugated with Alexa Fluor 594 and donkey anti-goat secondary antibody conjugated with Alexa Fluor 488 (Life Technologies) .	AB5405

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq