Document Identifier: **PMC8487208**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Our animal protocol was approved by the Institutional Animal Care and Use Committee at Sloan Kettering Institute.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Mice Female C57BL6J mice were purchased from the Jackson Laboratory (Stock 000664).
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
In all experiments, animals were randomly assigned to various experimental groups.
Blinding
not detected.
Power Analysis
The numbers of animals included in the study are discussed in each figure legend and were strictly limited by the necessary statistical power .
Replication
Quantitative real-time PCR was performed in triplicate with SYBR Green PCR Master Mix (Life Technologies) and an Applied Biosystems 7500 Real-time PCR Instrument (Life Technologies) , using gene-specific primers .

Number: Quantitative real-time PCR was performed in triplicate with SYBR Green PCR Master Mix (Life Technologies) and an Applied Biosystems 7500 Real-time PCR Instrument (Life Technologies) , using gene-specific primers .
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Polypeptides were separated by 15 SDS-PAGE , and western blot analysis was performed to determine the expression of mGM-CSF , using an anti-mGM-CSF antibody (Thermo Fisher) , or to investigate the activation status of different components of the cGASSTING pathway using antibodies against TBK-1 , p-TBK-1 , IRF-3 , p-IRF-3	anti-mGM-CSF		
	TBK-1 , p-TBK-1 , IRF-3 , p-IRF-3		
	p-TBK-1		
	IRF-3		
Cells were washed , permeabilized with PermBuffer (BD Biosciences) for 30 min on ice and stained with antibodies against CD11c and CD11b for 30 min .	antibodies against CD11c		
Cells were washed , permeabilized with PermBuffer (BD Biosciences) for 30 min on ice , and stained with antibodies against MHC Class I , CD40 , CD86 , and CD80 for 30 min .	antibodies against MHC Class I		
	CD40		
	CD86		
	CD80		
Flow cytometry analysis For tumor-infiltrating lymphocytes analysis , single cell suspensions were generated and processed for surface labeling with anti-CD3 , anti-CD45 , anti-CD4 , and anti-CD8 antibodies .	anti-CD3		
	anti-CD45		
	anti-CD4		
	anti-CD8		
Twenty-four hpi , cells were trypsinized and stained using an anti-PD-L1 antibody (eBioscience) .	anti-PD-L1		
The tumors were injected two times a week , concurrently with i.p. delivery of anti-CTLA-4 (100 ug per mouse) or isotype control antibodies .	anti-CTLA-4		
Murine anti-GM-CSF antibody was purchased from Thermo Fisher .	anti-GM-CSF		
Experimental Models: Cell Lines			
BSC-40 cells were maintained in DMEM medium containing 5 FBS, penicillin, and streptomycin.	BSC-40		Suggestion: RRID:CVCL_3656(link)
For cytokine assays , BMDCs or B16-F10 melanoma cells were infected with various viruses at an MOI of 10 for 1 hour , or mock-infected.	B16-F10		Suggestion: RRID:CVCL_0159(link)
Experimental Models: Organisms/Strains			
Batf3-- mice were from Dr Kenneth Murphy (Washington University).	Batf3--		

RNA isolation and quantitative real-time PCR B16-F10 melanoma cells were implanted into the right and left flanks of C57BL6J mice (5x105 cells into the right flank and 2.5x105 cells into the left flank) .	C57BL6J		Suggestion: RRID:IMSR_JAX:000664(link)
Recombinant DNA			
Generating recombinant vaccinia virus expressing mGM-CSF The murine GM-CSF (mGM-CSF) coding sequence was inserted into the pCB vector between Xba I and EcoR I sites.	pCB		Suggestion: RRID:Addgene_162699(link)
Software and Algorithms			
Data were analyzed using FlowJo software (Treestar)	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistics Two-tailed unpaired Students t-tests were used for comparisons of two groups in the studies .	Students t-tests
Survival data were analyzed by log-rank (Mantel-Cox) test .	log-rank
	Mantel-Cox) test
Oligonucleotides	
Primer sequences used for the PCR reactions were: TK-F2: 5-TGTGAAGACGATAAATTAATGA TC-3 pCB-R3: 5-ACCTGATGGATAAAAAGGCG-3 TK-F4: 5-TTGTCATCATGAACGGCGGA-3 TK-R4: 5-TCCTTCGTTTGCCATACGCT-3 GM-F: 5-GGCATTGTGGTCTACAGCCT-3 GM-R: 5-GTGTTTCACAGTCCGTTTCCG-3 TK-F5: 5-GAACGGGACTATGGACGCAT-3 TK-R5:	5-TGTGAAGACGATAAATTAATGA TC-3 pCB-R3: 5-ACCTGATGGATAAAAAGGCG-3 TK-F4: 5-TTGTCATCATGAACGGCGGA-3 TK-R4: 5-TCCTTCGTTTGCCATACGCT-3 GM-F: 5-GGCATTGTGGTCTACAGCCT-3 GM-R
	5-ACCTGATGGATAAAAAGGCG-3
	5-TTGTCATCATGAACGGCGGA-3
	5-TCCTTCGTTTGCCATACGCT-3
	GM-F
	5-GGCATTGTGGTCTACAGCCT-3
	5-GAACGGGACTATGGACGCAT-3
The primer sequences for quantitative real-time PCR were: Ifnb-F: 5-TGGAGATGACGGAGAAGATG-3 Ifnb-R: 5-TTGGATGGCAAAGGCAGT-3 Il6-F: 5-AGGCATAACGCACTAGGTTT-3 II6-R: 5-AGCTGGAGTCACAGAAGGAG-3 Ccl4-F: 5-GCCCTCTCTCTCCTCTTGCT-3 Ccl4-R: 5-CTGGTCTCATAGTAATCCATC-3 Ccl5-F: 5-GCCCACGTCAAGGAGTATTTCT A-3 Ccl5-R: 5-ACACACTTGGCGGTTCCTTC-3 Cxcl9-F: 5-GGAACCCTAGTGATAAGGAATG CA-3 Cxcl9-R: 5-TGAGGTCTTTGAGGGATTTGTA GTG-3 Cxcl10-F: 5-GTCAGGTTGCCTCTGTCTCA-3 Cxcl10-R: 5-TCAGGGAAGAGTCTGGAAAG-3 GAPDH-F: 5-ATCAAGAAGGTGGTGAAGCA-3 GAPDH-R: 5-AGACAACCTGGTCCTCAGTGT- 3 Unilateral intradermal tumor implantation and intratumoral injection of viruses B16-F10 melanoma cells (1x105 cells in a volume of 50 uL PBS) were implanted intradermally into the shaved skin on the right flanks of WT C57BL6J or Batf3--mice.	5-TGGAGATGACGGAGAAGATG-3
	5-TTGGATGGCAAAGGCAGT-3
	5-AGGCATAACGCACTAGGTTT-3
	5-AGCTGGAGTCACAGAAGGAG-3
	5-GCCCTCTCTCTCCTCTTGCT-3
	5-CTGGTCTCATAGTAATCCATC- 3
	5-GCCCACGTCAAGGAGTATTTCT A-3
	5-ACACACTTGGCGGTTCCTTC-3
	5-GGAACCCTAGTGATAAGGAATG CA-3
	5-TGAGGTCTTTGAGGGATTTGTA GTG-3
	5-GTCAGGTTGCCTCTGTCTCA-3
	5-TCAGGGAAGAGTCTGGAAAG-3
	5-ATCAAGAAGGTGGTGAAGCA-3 GAPDH-R
Unresolved RRIDs	

E3L83N (OV-TK+), OV, or OV-GM viruses were propagated in BSC-40 (Africa green monkey kidney cells, ATCC-CRL2761).	CRL2761

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq