Document Identifier: **PMC8489026**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Field Sample Permit: Table 1 Natural pea aphid lineages considered in the present study Name Sampling location Collection date Biotype Secondary symbiont Gt1 Bugey (France) June 2014 Genista tinctoria Hamiltonella defensa Gt2 Bugey (France) June 2014 G
Inclusion and Exclusion Criteria
This latter was estimated by dividing the number of aphid mummies (i.e. , dead aphids containing a parasitoid immature) by the total number of aphids successfully attacked transferred onto the host plant and recovered at that time (i.e. dead or missing individuals were excluded from analyses) .
Attrition
not required.
Sex as a biological variable
These six natural aphid lineages came from single aphid females sampled in the field on their respective host plants (details in Table 1) .
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
Bacterial secondary symbionts are not randomly distributed among pea aphid biotypes but instead form preferential associations with some biotypes (Henry et al. 2013 Mathe-Hubert et al. 2019) .
Blinding
not detected.
Power Analysis
not detected.
Replication

In this model , the symbiotic status of aphids was fit as fixed factor along with the tibia size of the parasitoid mother as covariate , whereas the replicate ID was fit as random factor (i.e. , some parasitoid offspring developed on the same aphid host plant and environmental conditions) .

Cell Line Authentication

Authentication: They were assigned to their corresponding biotype by genotyping and comparing their microsatellite profile to a reference database (Peccoud et al. 2009 Peccoud et al. 2015) .

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains			
Transfer of some symbionts in new aphid hosts did not work (Sochard et al. 2020) and a few lineages were lost between their creation and the experiments (Gt2-HdMs1 , Gt2-HdMs2 and Ms2-HdGt2) , so we could not test all possible combinations .	Gt2-HdMs2		
Ms1-Ri T2 Ms1 New R . insecticola T2 Ms2-Hd- Ms2 Cured No SS - Ms2-Hd Ms2 Ms2 Control H . defensa Ms2	SS - Ms2-Hd Ms2 Ms2 Control H		
Software and Algorithms			
The mass rearing of A . ervi started from at least 50 males and females produced by Koppert Biological Systems(c) .	Koppert Biological Systems(c		

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq