Document Identifier: **PMC8490173**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: The contents of the five educational sessions were as follows: the first session included an introduction of the study and its objectives, obtaining patient consent, and filling the questionnaires.
Inclusion and Exclusion Criteria
Patients who were known to have a terminal or severe illness or mental impairment were excluded. 3. Randomization We had a sample frame consisting of a list of all clinic attendants who fulfilled the inclusion criteria.
Attrition
not detected.
Sex as a biological variable
Current (smoking almost every day in the past 3 months) smokers of both sexes aged between and 18-64 years who are visiting the family medicine clinic at Cairo University Hospitals for a variety of medical conditions (e.g., diabetes mellitus and hypertension) were invited to participate in the study.
Subject Demographics
Age: Current (smoking almost every day in the past 3 months) smokers of both sexes aged between and 18-64 years who are visiting the family medicine clinic at Cairo University Hospitals for a variety of medical conditions (e.g., diabetes mellitus and hypertension) were invited to participate in the study.
Randomization
Setting This was a single-blinded randomized controlled clinical trial that was conducted on 78 current smokers visiting family medicine outpatient clinics at Cairo University Hospitals in Egypt.
Blinding
not detected.
Power Analysis
Participants The sample size was calculated using the Epi 6 program set (Centers for Disease Control and Prevention, Atlanta, GA, USA) to achieve 100 power and to detect a difference of 13.0 (5 reduction of nicotine dependence scale) between the null hypothesis (wherein both groups means are 259.0) and the alternative

hypothesis (wherein the mean of group 2 is 246.0), with estimated group standard deviations of 1.0, and with a significance level (a) of 0.05000 using a two-sided two-sample t-test 11.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The total score ranged from 37 to 259, with higher scores indicating greater levels of nicotine dependence. 6. Statistical Analysis of Data Data analysis was performed using the IBM SPSS ver. 21.0 (IBM Corp., Armonk, NY, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Participants The sample size was calculated using the Epi 6 program set (Centers for Disease Control and Prevention, Atlanta, GA, USA) to achieve 100 power and to detect a difference of 13.0 (5 reduction of nicotine dependence scale) between the null hypothesis (wherein both groups means are 259.0) and the alternative hypothesis (wherein the mean of group 2 is 246.0), with estimated group standard deviations of 1.0, and with a significance level (a) of 0.05000 using a two-sided two-sample t-test 11.	t-test
A chi-square test was used to compare categorical data across groups and in repeated measures, while Fishers exact test was used to compare qualitative variables between groups when one expected cell or more had a value less than five.	chi-square test
	Fishers exact test
Pairwise testing was performed using paired and independent t-tests with Bonferroni adjustment for the number of comparisons.	t-tests
	Bonferroni
Mann-Whitney and Wilcoxon tests were used instead of unpaired t-test and paired t-test, respectively, to analyze non-parametric data if the standard deviation was more than 30 of the mean.	Mann-Whitney
	Wilcoxon tests

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq