Document Identifier: **PMC8494515**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: Written informed consent was obtained from each participant andor their parentslegal guardians (as appropriate) local research ethics boards approved the study .
IRB: Written informed consent was obtained from each participant andor their parentslegal guardians (as appropriate) local research ethics boards approved the study .
Inclusion and Exclusion Criteria
Inclusion criteria included a confirmed diagnosis of severe hemophilia A , defined in this study as a baseline FVIII:C level of <2 participant receiving ADVATE for prophylaxis and body weight >=12 kg and <=120 kg .
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Weight: Inclusion criteria included a confirmed diagnosis of severe hemophilia A , defined in this study as a baseline FVIII:C level of <2 participant receiving ADVATE for prophylaxis and body weight >=12 kg and <=120 kg .
Age: The myPKFiT dosing tool was built from the ADVATE pre-licensure PK data using 184 full PK datasets (10 postinfusion samples) from 100 adultsadolescents with hemophilia A (10-65 years of age) and 52 reduced sample PK datasets (four postinfusion samples) from 52 boys with hemophilia A (1-6 years of age) .
Randomization
not detected.
Blinding
not detected.
Power Analysis

A sample size of 35 patients was deemed necessary to achieve 80 power to detect an ICC of 0.8 under the alternative hypothesis, when the ICC under the null hypothesis is 0.5 using an F-test with a significance level of 0.01.	
Replication	
not required.	
Protocol Information	
Identifiers: The study was registered on ClinicalTrials . gov (NCT02750085) .	NCT02750085

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
VWF:Ag levels were measured by enzyme-linked immunosorbent assay (ELISA) using polyclonal rabbit anti-human VWF-coating antibody (Ab) (A0082 Dako , Glostrup , Denmark) ,	anti-human VWF-coating		
	A0082		
VWF propeptide (VWF:pp) levels were measured by ELISA using Ab pair anti-human VWF:pp MW1939 (Antibody Chain , Utrecht , Netherlands) .	anti-human VWF:pp MW1939		
Experimental Models: Organisms/Strains			
Pharmacokinetic Parameters and Analysis The PK parameters reported herein were chosen in accordance with the guidelines from the FVIII and FIX Subcommittee of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis (ISTH) .	FVIII		Suggestion: RRID:NSRRC_0018(link)
Software and Algorithms			
14 PK parameters from the 6-point PK were calculated using Phoenix WinNonlin 7.0 (Certara USA , Inc . Princeton , United States) , considered the gold standard for analyzing individual PK data , and ADVATE myPKFiT v .	Phoenix WinNonlin		
The model and its development are detailed in Bjorkman et al. 11 Statistical Analyses Statistical analyses were conducted using MedCalc Statistical Software version 18.10.2	MedCalc		Suggestion: MedCalc (RRID:SCR_015044)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
17 18 Pearson correlation coefficients were used to determine the extent of linear correlation between PK parameters and (1) VWF:Ag (2) VWF:pp (3) VWF:ppAg ratio and (4) age.	Pearson correlation
19 Paired t -tests were used to compare FVIII:C levels from the one-stage and chromogenic FVIII:C assays, and one-way analysis of variance (ANOVA) was used for comparing VWF:Ag levels in O versus non-O blood group subjects.	t -tests
	analysis of variance
	ANOVA
A sample size of 35 patients was deemed necessary to achieve 80 power to detect an ICC of 0.8 under the alternative hypothesis, when the ICC under the null hypothesis is 0.5 using an F-test with a significance level of 0.01.	F-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq