Document Identifier: **PMC8496413**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Approval was obtained from the University of Michigan Institutional Review Board and Aravind Madurai Ethics Committee.
Inclusion and Exclusion Criteria
Inclusion criteria were patients 15 years or older with MK as diagnosed by the treating clinician.
Attrition
not detected.
Sex as a biological variable
Patients were excluded from participation if from a vulnerable population (i.e., hospitalized, imprisoned, institutionalized) or pregnant or not able to provide consent.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
For each algorithm, the participants were randomly divided into six groups of approximately equal size.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Manual Segmentation Manual segmentation of the ROIs on the SLP images was performed by ophthalmologists (MFK, KHK) using ImageJ software (National Institutes of Health, Bethesda, MD, USA), as described in prior work.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Implementation The data set was prepared using ImageJ and MATLAB (MathWorks, Natick, MA, USA).	MATLAB		Suggestion: MATLAB, RRID:SCR_001622(link)
The automatic segmentation and BCVA estimation algorithms were implemented in Python using the TensorFlow 35 (version 1.5.1), Keras 36 (version 2.0.8), and Scikit-learn 37 (version 0.22.2) libraries.	Python		Suggestion: python-bx (RRID:SCR_024202)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Correlation With BCVA Pearsons correlation was calculated to analyze the relationship between the measurements of pathologic ROIs obtained from either manual or automatic segmentations and BCVA.	Pearsons correlation
Statistical Significance Tests To determine the statistical significance of the difference between using manual and automatic segmentations, Williams test was used for the differences in correlation, and the paired-sample t-test was used for the differences in absolute error.	Williams test
	t-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq