Document Identifier: **PMC8502538**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not detected.
Inclusion and Exclusion Criteria
Patients on antibiotics or nonsteroidal anti-inflammatory drugs within 4 weeks prior to OGDS and those with gastric malignancy or immunosuppression were excluded.
Attrition
Three biopsies were obtained from antrum andor corpus of the stomach from each patient.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Samples were run in duplicate and were considered positive if both reactions were positive.
Number: Samples were run in duplicate and were considered positive if both reactions were positive.
Cell Line Authentication

Authentication: Detection of Point Mutation in the 23S rRNA Gene of H. pylori in Biopsy Samples by Multiplex qPCR The ureA-positive samples were subjected to multiplex qPCR for the detection of mutation in 23S rRNA gene.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
Among the 59 H. pylori isolates H. pylori strains A3, A26, C81, C55, A142, A129, A150, A177, and C192 were used as positive controls.	C81		
Experimental Models: Organisms/Strains			
33 The limit of detection of the ureA qPCR assay was determined using 10 serially diluted DNA samples of H. pylori strain C55.	C55		
The specificity of the 23S rRNA gene qPCR assay was determined by testing other bacterial strains including Acinetobacter spp., B. subtilis, S. flexneri, P. mirabilis, Serratia spp, S. typhimurium, E. coli, S. epidermidis, P. aeruginosa, Enterobacter spp., K. pneumoniae, V. parahaemolyticus, V. cholera, S. aureus, and Flavobacterium spp. H. pylori ATCC 43526 and H. pylori J99 were used as positive controls.	P. aeruginosa		
Software and Algorithms			
The published sequences of H. pylori 23S rRNA gene (Accession no. KC556778), gyrA (Accession no. L29481), the wild type H. pylori ATCC 26695, and sequences from clinical H. pylori strains were aligned using CLUSTAL W Multiple Alignment30 in Bioedit Version 7.2.5, and primers and probes were designed by using the online PrimerQuest Tool on the IDT website (https:sg.idtdna.comPrimerQuestHomeIndex).	Bioedit		Suggestion: BioEdit (RRID:SCR_007361)(link)
The primers and probes were synthesized by Integrated DNA Technologies (Singapore Science Park II, Singapore) and Macrogen (Synapase, Singapore) and are listed in Table 1.	Macrogen		Suggestion: Macrogen (RRID:SCR_014454)(link)
After sequencing, a BLAST search was performed to check the sequence similarityidentity of the DNA sequences.	BLAST		Suggestion: WU-BLAST (RRID:SCR_011824)(link)
Statistical Analysis Statistical analysis was performed using SPSS software version 23 (SPSS Inc, Chicago, IL, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Differences between groups of the categorical data were statistically evaluated using Pearson Chi-square (kh2) test.	Pearson Chi-square
The independent samples t-test is used to determine the positivity results between singleplex and multiplex qPCR.	t-test
Differences of qPCR-negative and qPCR-positive were statistically evaluated using one sample Chi-square (kh2) test.	Chi-square
Kappa and McNemar tests were used to compare consistency between different detection methods.	McNemar tests
Oligonucleotides	
Table 1 Oligonucleotides Sequence Used in This Study PrimersProbes Sequences (5-3) Targets Product (bp) UreA-Fa AGTTCCTGGTGAGTTGTTCTT ureA 120 UreA-Rb TGGAAGTGTGAGCCGATTT Cla-Fa CTGCGCATGATATTCCCGTT 23S rDNA 157 Cla-Rb CGACCTGCATGATGGCGTA Lev-Fa CGCCATCAATAGAGCCAAAG gyrA 136 Lev-Rb GTGGGTGATGTGATTGGTAAAT UreA-Pc FAM-AAATGTTGGCGACAGACGGT TC-TAMRA ureA Cla-PWTc Cy5-AAGGTCCACGGGGTCTTTCC GTC-BHQ2 Wild type Cla-P2142Gc HEX-AGGTCCACGGGGTCTTCCCG TCT-BHQ2 A2142G Cla-P2143Gc FAM-AGGTCCACGGGGTCTCTCCG TCTT-BHQ2 A2143G Lev-N87WTc HEX-CCG +C+G+TTA+T+CG-IowaBlack(r)FQ Wild type Lev-N87Kc 6-FAM-CCG+C+T +TT+A+T+CGC-IowaBlack(r)FQ N87K Lev-N87Ic Cy5-CCG+C+A+AT+A+T +CGC-IowaBlack(r)RQ-Sp N87I Lev-D91WTc HEX-TA+T+G+AT+GCG+CT-IowaBlack(r)FQ Wild type Lev-D91Nc Cy5-TTA+T+A+ATG+C+G+CTAGT-IowaBlack(r)RQ-Sp D91N Lev-D91Yc 6-FAM-TTA+T+T+ATG+C+G+CTAGT-IowaBlack(r)FQ D91Y Notes: aForward primer, bReverse primer, cTaqMan probe.	TGGAAGTGTGAGCCGATTT CGACCTGCATGATGGCGTA GTGGGTGATGTGATTGGTAAAT

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq