Document Identifier: **PMC8503744**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Animals A total of 12 male BALB/c nude mice aged 4 weeks (weight, 16.86±9.3 g) for the in vivo study were provided by Beijing Vital River Laboratory Animal Technology Co., Ltd., and reared according to the Animal Research Committee's guidelines of Anhui Medical University (Hefei, China).
Euthanasia Agents: At day 24 after RO injection, the mice were anesthetized with an intraperitoneal injection of pentobarbital (50 mg/kg) and euthanized via cervical dislocation.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: Animals A total of 12 male BALB/c nude mice aged 4 weeks (weight, 16.86±9.3 g) for the in vivo study were provided by Beijing Vital River Laboratory Animal Technology Co., Ltd., and reared according to the Animal Research Committee's guidelines of Anhui Medical University (Hefei, China).
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication

Following trypsinization , ~2,000 PANC-1 cells in the logarithmic growth phase were prepared as a unicellular suspension to seed in a 96-well cell culture plate with 100 µl volume in triplicate.
Number: Following trypsinization , ~2,000 PANC-1 cells in the logarithmic growth phase were prepared as a unicellular suspension to seed in a 96-well cell culture plate with 100 µl volume in triplicate.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Specific antibodies against β-actin, p27, phosphorylated (p)ERK1/2, ERK1/2, pJNK, JNK, cyclin B1 and cyclin E were purchased from Santa Cruz Biotechnology, Inc.	p27, phosphorylated (p)ERK1/2, ERK1/2, pJNK, JNK		
	cyclin B1		
	cyclin E		
After non-specific blocking in 5 % skimmed milk for 2 h at room temperature , the PVDF membranes were washed with PBS containing 0.1 % Tween-20 three times and incubated at 4°C overnight with specific primary antibodies against β-actin (cat. no. sc-47778; 1:1,000)	β-actin		
Subsequently , the membranes were reacted with the corresponding HRP-conjugated secondary antibody (goat anti-mouse IgG; cat. no. AP130P; 1:10,000) diluted in TBS solution for 2 h at room temperature.	anti-mouse IgG		
For immunohistochemical staining, the sections(4-μm thick) obtained from the formalin-fixed and paraffin-embedded tissues were blocked with 3% BSA (Beyotime Institute of Biotechnology) diluted in 0.2% Triton X-100/PBS at room temperature for 30 min and stained with a Ki67 specific antibody (cat. no. ZA-0502; OriGene Technologies, Inc.) overnight at 4°C, followed by incubation with HRP-conjugated secondary antibody (cat. no. PV-6001) for 2 h at room temperature and development with a DAB Horseradish Peroxidase Color Development kit (cat. no. ZLI-9018; OriGene Technologies, Inc.).	Ki67		
Experimental Models: Cell Lines			
Following trypsinization , ~2,000 PANC-1 cells in the logarithmic growth phase were prepared as a unicellular suspension to seed in a 96-well cell culture plate with 100 μl volume in triplicate.	PANC-1		
Experimental Models: Organisms/Strains			

Animals A total of 12 male BALB/c nude mice aged 4 weeks (weight, 16.86±9.3 g) for the in vivo study were provided by Beijing Vital River Laboratory Animal Technology Co., Ltd., and reared according to the Animal Research Committee's guidelines of Anhui Medical University (Hefei, China).	BALB/c nude		
Software and Algorithms			
Data analysis was performed using FlowJo software (FlowJo version 10.5.4; FlowJo LLC) .	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)
The signals of the specific bands were identified and detected with the Quantity One analysis software (version 4.6.6; Bio-Rad Laboratories , Inc .)	Quantity One		
	Bio-Rad Laboratories		
Statistical analysis All data were analyzed using SPSS version 15.0 software (SPSS, Inc.).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The significant differences between groups were determined using one-way ANOVA followed by a Tukey's post hoc test.	ANOVA
	Tukey's post hoc test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq