Document Identifier: **PMC8503845**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not required.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Number: Three times washing with native lysis buffer were applied to protein A beads , and substrates were eluted with 1X SDS sample buffer
Data Information
Availability: Materials Availability All uniquestable reagents generated in this study are available from the Lead Contact without restriction.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Samples were then resolved on 10 SDS-PAGE and subject to western blot analysis using the indicated antibodies or antisera.	antisera		
Cell lysis was conducted by agitation with glass beads, and the indicated GFP tagged proteins were immunoprecipitated with a 2 h incubation with anti-GFP antibody (Invitrogen) followed by overnight incubation with protein A Sepharose beads (GE healthcare).	anti-GFP		
Polyubiquitin and the precipitated substrate were visualized using anti-ubiquitin (P4D1) anti-HA (3F10) antibodies , respectively .	anti-ubiquitin		
	3F10		
The supernatant was next transferred to new micro-centrifuge tube with HA antibody (12CA5 Roche) for 4 hours incubation at 4degC .	HA		
Recombinant DNA			
Both SZ*-GFP and the Ub4 fragments were then inserted into p416TEF plasmid digested with XbaI and HindIII, thus generating plasmid pSZ10.	p416TEF		Suggestion: RRID:Addgene_99553(link)
Plasmid pSZ11 that expresses Ydj1-HPDQAA was generated by site-directed mutagenesis of BPM390 using primers OSZ17 and OSZ18.	pSZ11		
To create the pSZ12, SZ*-GFP-Ub4* construct, we first ordered the Ub4* sequence from IDT with the L8E, I44E, and V70D mutations (on top of the K29R, K48R, and K63R mutations that are localized in SZ*-GFP-Ub4 construct) with BamHI and XhoI sites on the N- and C-terminal, respectively.	pSZ12		
For plasmid pSZ13, XbaI and XhoI enzymes were used to cut plasmid pSZ03 to release SZ*-GFP fragment, which was then ligated with 415TEF plasmid treated with same enzymes.	pSZ13		
	pSZ03		
In order to generate plasmid pSZ14, which harbors ED-Ub4, the ED-Wsc1-L63R fragment was first amplified using primers OSZ21 and OSZ22 and pSW144 as a template.	pSZ14		
	pSW144		

The vector containing the Ub4 fragment was made by cutting SZ*-GFP fragment from pSZ10 using XbaI and BamHI.	pSZ10		
Yor1-GFP-Ub4 (pSZ15) and Pca1(1-392)-Yor1-GFP-Ub4 (pSZ16) were constructed by Gibson assembly.	pSZ15		
	pSZ16		
COPII protein purification Sar1 was purified from the pTY40 plasmid (Barlowe et al. , 1994) in an E . coli BL21-based CBB205 strain (NEB) .	pTY40		
Software and Algorithms			
METHOD DETAILS Yeast strains and plasmid construction Plasmids and primers used in this study are listed in the STAR Methods.	STAR		
The DNA sequences of all constructs were confirmed by Genewiz.	Genewiz		
Band intensities from western blotting and pulse chase experiments were quantified using ImageJ version 1.48V and normalized to the loading control or to total protein.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
GFP fluorescence intensities were quantified using FIJI (2.0.0-rc-691.52i).	FIJI		
Statistical analysis in this work was determined by Prism 7 using either unpaired t- tests, ordinary one-way ANOVA with Holm-Sidaks multiple comparison tests, or ordinary two-way ANOVA with a Dunnetts multiple comparisons test.	Prism		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis in this work was determined by Prism 7 using either unpaired t- tests, ordinary one-way ANOVA with Holm-Sidaks multiple comparison tests, or ordinary two-way ANOVA with a Dunnetts multiple comparisons test.	ANOVA
	Holm-Sidaks multiple comparison tests
	Dunnetts multiple comparisons test
Unresolved RRIDs	
The Ub4 fragment was generated and collected by cutting plasmid 1677 with BamHI and HindIII.	plasmid 1677
In brief, a vector fragment was generated by digesting plasmid 105 with XbaI and XhoI enzymes.	plasmid 105

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq