Document Identifier: **PMC8504217**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: All participants provided written informed consent underwent a standardized history, anthropometric measurements, physical examination, and biochemical testing.
IRB: 34 This study approved by the Institutional Review Board at the University of California, San Diego (UCSD) and the UCSD Clinical and Translational Research Institute.
Inclusion and Exclusion Criteria
Exclusion criteria for this study included any of the following: excessive alcohol use 14 (men) or 7 (women) drinks per week, use of steatogenic or hepatotoxic medications, and inadequate liver biopsy (biopsy length was <10 mm or portal triads <10).
Attrition
Five trained operators performed VCTE and 2D-SWE examinations, although the majority were performed by one operator (n = 65 participants).
Sex as a biological variable
Exclusion criteria for this study included any of the following: excessive alcohol use 14 (men) or 7 (women) drinks per week, use of steatogenic or hepatotoxic medications, and inadequate liver biopsy (biopsy length was <10 mm or portal triads <10).
Subject Demographics
Age: Methods Study population and liver histology This is a cross-sectional analysis of consecutive adults (18 years of age) with NAFLD enrolled in prospective cohort study who underwent 2D-SWE and VCTE on the same day at the University of California at San Diego.
Weight: Subgroups were defined according to BMI quartiles: patients with BMI < 34.3 kgm2 and patients with BMI 34.3 kgm2 (75th percentile).
Randomization
not detected.
Blinding

Liver biopsy was considered as the reference standard for liver fibrosis stage, and all liver biopsies were reviewed by one expert liver pathologist blinded to all clinical and histological data and scored using the Nonalcoholic Steatohepatitis Clinical Research Network (NASH CRN) histological scoring system.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis Categorical data were compared using the chi-square and Fishers exact tests.	chi-square
	Fishers exact tests
Distributions of continuous variables were analyzed using the Mann-Whitney U test.	Mann-Whitney U test
The association between liver stiffness and histological fibrosis stage was evaluated by Spearmans rank correlation test.	Spearmans rank correlation test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq