Document Identifier: **PMC8504813**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: METHODS Animal experiments All animal experiments were performed in accordance with the guidelines of the Guide for the Care and Use of Laboratory Animals and were approved by the Bioethics Committee of Second Military Medical University.
Field Sample Permit: After collection of the douche and determination of the embryonic development under the microscope (dropwise addition of the sample used for microscopic examination, 50 ul volume of each drop), we determined the number of 2-16-cell stage embryos, morula, and blastocysts, and calculated the total number of embryos in each mouse.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Four-week-old Institute of Cancer Research (ICR) female mice (mean weight +- SD: 16 +- 2.1 g) were purchased from the Shanghai Research Center for Model Organisms and were raised in the specific pathogen-free (SPF) animal facility of the Department of Cell Biology, SMMU, under a controlled environment (14-h light10-h dark cycle, 24degC).
Subject Demographics
Age: Four-week-old Institute of Cancer Research (ICR) female mice (mean weight +- SD: 16 +- 2.1 g) were purchased from the Shanghai Research Center for Model Organisms and were raised in the specific pathogen-free (SPF) animal facility of the Department of Cell Biology, SMMU, under a controlled environment (14-h light10-h dark cycle, 24degC).
Randomization
not detected.
Blinding
not detected.
Power Analysis

not detected.
Replication
Experimental design and statistical analysis Each experiment was repeated at least three times.
Number: Experimental design and statistical analysis Each experiment was repeated at least three times.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The bound antibodies were detected using HRP-conjugated anti-rabbit and peroxidase-conjugated anti-mouse antibodies, and analyzed using chemiluminescence (ECL System) via autoradiography.	anti-rabbit		
	anti-mouse		
Software and Algorithms			
Cell culture KGN cells (human ovarian granulosa cell line, GCs) were originally obtained from the Cell Bank at the Shenzhen Institute, Chinese Academy of Sciences (Shenzhen, Peoples Republic of China) and cultured in DMEM medium (Gibco, C11995500BT) containing 10 fetal bovine serum (Gibco, 10091-148) and 1 100x penicillin-streptomycin solution (Gibco, 15140-122) in an incubator (37degC, 5 CO2, 95 humidity).	Gibco		
Statistical analysis was performed using the Students t-test with the SPSSGraphPad Prism software.	SPSSGraphPad Prism		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis was performed using the Students t-test with the SPSSGraphPad Prism software.	t-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq