Document Identifier: **PMC8506827**

# SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

**Table 1: Rigor Adherence Table**

| |
|---|
| Ethics |
| Field Sample Permit: Sampling of late-stage larvae and pupal nests were conducted during June and July, respectively, via timed observations. |
| Inclusion and Exclusion Criteria |
| not required. |
| Attrition |
| not required. |
| Sex as a biological variable |
| Materials and Methods Study Organism The browntail moth life cycle is initiated in the later part of July in Maine when females lay their eggs in masses on the undersides of leaves at the terminal ends of the branches in the upper canopy of host trees (Fig. 4). |
| Subject Demographics |
| Age: not required. |
| Weight: not required. |
| Randomization |
| Over two weeks, three randomly selected hibernacula from each site were removed from cold storage and placed in a Fabri-Kal deli cup container (473 ml plastic cup) or plastic freezer bag. |
| Blinding |
| not detected. |
| Power Analysis |
| not detected. |
| Replication |
| not required. |

**Table 2: Key Resources Table**

| Your Sentences | REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|---|---|---|
| Software and Algorithms | | | |
| The amount of potential habitat present at each site was determined by identifying a 1.5 km radius around the center of the site using ArcGIS version 10.1 ( ESRI , Redlands , CA) . | ArcGIS | | Suggestion: ArcGIS for Desktop Basic (RRID:SCR_011081)( link) |

**Other Entities Detected**

| Your Sentences | Recognized Entity |
|---|---|
| Statistical Tests | |
| Additionally , differences between years and sites and mean mortality of laboratory-reared larvae and pupae were examined in RStudio ( RStudio 2019 , version 1.1.414 ) using two-way analysis of variance on an untransformed dependent variable ( mean mortality ) as the errors were homogeneous and normally distributed. | analysis of variance |
| ± 1.597 0.151 ± 0.073 0.248 ± 0.065 x 2 ( 1 ) = 2.769 , P = 0.096 x 2 ( 1 ) = 4.282 , P = 0.039 x 2 ( 1 ) = 14.632 , P < 0.001 *Likelihood ratio test , Chi-square test for difference between whole ( all independent variables ) and reduced ( intercept only ) model . **To predict life-stage relative abundance models fit with Poisson or negative binomial exponentiate the linear additive equation of model coefficients . | Chi-square test |

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq