Document Identifier: **PMC8517348**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Euthanasia Agents: Cell culture and treatment Ventricular mouse cardiomyocytes were isolated by a standard enzymatic technique as described before. 16 Briefly , after cervical dislocation , hearts were rapidly removed and perfused at 37degC for 10 min with a bicarbonate buffer (pH 7.4) containing low Ca2+ and collagenase (type II PAN BIOTECH) .
Inclusion and Exclusion Criteria
Exclusion criterion was defined as the mouses inability to continue regular treadmill running and to maintain the required running speed and distance despite mechanical prodding
Attrition
28 TA muscle from aged adult male Ahnak1-KO (n = 16), male WT (n = 10), female Ahnak1-KO (n = 11), and female WT (n = 7) mice was dissected, transferred to an organ bath containing Krebs solution (vol. 20 mL) bubbled with 95 O2 and 5 CO2 and kept at constant temperature (22degC) where it was mounted between the hook of a force transducer (FT-03 AstroNova) and a hook connected to a movable shaft that was used to adjust the muscle length.
Sex as a biological variable
Material and methods Animals Fifteen- to 16-month-old male and female homozygous Ahnak1-KO 10 and their WT littermates were used.
Subject Demographics
Age: Material and methods Animals Fifteen- to 16-month-old male and female homozygous Ahnak1-KO 10 and their WT littermates were used.
Randomization
not detected.
Blinding
For analyses of mitochondrial size (surface area), 10 micrographs were taken for each LV and TA tissue to measure mitochondrial size per unit area (48 mm2) in a blinded fashion.
Power Analysis
not detected.

Replication
Twelve replicates were run for each condition , and experiments were repeated seven times independently .
Number: Twelve replicates were run for each condition , and experiments were repeated seven times independently .
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Labelling was performed with a primary anti-Ahnak1 antibody (E-5, Santa Cruz) for 1 h at 4degC and a secondary anti-mouse 12 nm colloidal gold conjugate (115-205-146, Dianova) for 30 min at RT.	anti-Ahnak1		
	E-5		
	anti-mouse		
Experimental Models: Cell Lines			
For in vitro investigations , we used a human adult left ventricular cardiomyocyte cell line , the AC16 cells , 17 which were cultured as previously described.	AC16		Suggestion: RRID:CVCL_4U18(link)
Experimental Models: Organisms/Strains			
Whole murine heart from aged adult Ahnak1-KO and WT mice of both sexes (n = 6 per group) were dissected and externally perfused with 4 (wv) formaldehyde0.5 (vv) glutaraldehyde in 0.1 M phosphate buffer.	Ahnak1-KO		
Software and Algorithms			
Band intensities were quantified with Image Lab software 5.1 V (Bio-Rad Laboratories) .	Image Lab		Suggestion: Image Lab Software (RRID:SCR_014210)(link)
	Bio-Rad Laboratories		Suggestion: Bio-Rad Laboratories (RRID:SCR_008426)(link)
Statistical analysis was performed using one-way analysis of variance followed by Bonferroni post hoc test to compare multiple groups, two-way analysis of variance followed by Bonferroni post hoc test to compare means, two-tailed Students t-test to compare values between two groups, or multiple t-test, as appropriate, using GraphPad Prism 5.01.	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis was performed using one-way analysis of variance followed by Bonferroni post hoc test to compare multiple groups, two-way analysis of variance followed by Bonferroni post hoc test to compare means, two-tailed Students t-test to compare values between two groups, or multiple t-test, as appropriate, using GraphPad Prism 5.01.	analysis of variance
	Bonferroni post hoc test
	t-test
	multiple t-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq