Document Identifier: **PMC8519618**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: All participants signed informed consent forms.
Inclusion and Exclusion Criteria
The exclusion criteria included CVD (except CSX), diabetes, any endothelial dysfunction diseases, asthma, pulmonary-renal disorders, active infection, and malignant disease.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: Materials and Methods The present study was conducted on 40 patients with CSX (51.7 ± 10.7 years of age) referring to the Department of Cardiology of Urmia University of Medical Sciencesý, Urmia, Iran, due to chest pain.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The MP concentration of the PPP samples was quantitatively determined through flow cytometry equipped with a forward scatter (FSC) photomultiplier tube , using specific anti-human CD31-FITC , CD41a-PE , CD62E-PE , and CD144-PE monoclonal antibodies (eBioscience Inc . , San Diego , CA , USA) .	anti-human CD31-FITC		
	CD144-PE		
Then , 300 µl of PPP was added to 3 special tubes (D-51588; SARSTEDT AG & Co . KG , Germany) equally (100 µl per tube) , and each tube was separately incubated in the dark with either CD144-PE , CD62E-PE , or both CD31-FITC and CD41a-PE antibodies at 4 °C for 45 minutes .	CD62E-PE		
	CD41a-PE		
Two distinct tubes containing anti-mouse PE-conjugated and FITC-conjugated IgG1 antibodies (eBioscience Inc . , San Diego , CA , USA) were employed as isotype controls in parallel with the test .	anti-mouse		
	FITC-conjugated IgG1		
Software and Algorithms			
cholesterol , triglyceride (TG) , high-density lipoprotein (HDL)-c , and low-density lipoprotein (LDL)-c level measurements were performed using commercially available kits (Pars Azmoon Co . , Tehran , Iran) in an automated analyzer (model ALCYON 300; Abbott Laboratories , Chicago , IL , USA) .	Abbott Laboratories		
Data were analyzed using SPSS software (version 19; SPSS Co., Chicago, IL, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
First, the Kolmogorov-Smirnov normality test was performed.	Kolmogorov-Smirnov normality test
The age effect, as a confounding factor, was checked for each quantitative variable, and mean of values was compared between patient and healthy groups using independent t-test.	t-test
Nominal variables were examined using chi-square or Fisher's exact tests.	chi-square
	Fisher's exact tests
The association between the variables was detected using the Pearson and Spearman correlation tests.	Pearson
	Spearman correlation

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq