Document Identifier: **PMC8552833**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
Eleven operated lizards were utilized after 11 (n = 3), 19 (n = 2), 22 (n = 3), and 34 (n = 3) days post-injury.
Sex as a biological variable
Briefly, adult lizards of both sexes were utilized, and the surgical procedures utilized in the study followed the regulations on animal care and experimental procedures under the Italian Guidelines (art. 5, DL 11692).
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
After western blotting, membranes were stained with Ponceau red to verify the protein transfer and incubated with a primary anti-mouse antibody for GAP-43 or an anti-rabbit antibody for nerurofilament-200 (see below).	GAP-43		
	anti-rabbit		
After extensive rinsing, the nitrocellulose strip was incubated with secondary FITC-conjugated antibodies (anti-mouse or anti-rabbit) diluted 1:2000 for 1 hour.	anti-mouse		
A mouse monoclonal antibody (G-9264 from Sigma) was utilized to detect GAP-43 (or neuroregulin) , a marker of axonal growth cones and nerve cells .	neuroregulin		
After repeated rinses in buffer , the sections were incubated with goat anti-mouse or anti-rabbit secondary antibodies conjugated to fluorescein isothyocyanate (FITC , Sigma , producing a green fluorescence) or tetramethyl rhodamine iso-thiocyanate (TRITC , Sigma , producing a red fluorescence) , mounted in anti-fading Fluoromount (Sigma) , and observed under a fluorescent microscope.	anti-fading Fluoromount (Sigma) ,		
Software and Algorithms			
After repeated rinses in buffer , the sections were incubated with goat anti-mouse or anti-rabbit secondary antibodies conjugated to fluorescein isothyocyanate (FITC , Sigma , producing a green fluorescence) or tetramethyl rhodamine iso-thiocyanate (TRITC , Sigma , producing a red fluorescence) , mounted in anti-fading Fluoromount (Sigma) , and observed under a fluorescent microscope.	Sigma		Suggestion: sigma-tau (RRID:SCR_000488)(link)
The sections were studied under a fluorescence microscope , and images were recorded by a digital camera and imported into a computer for making figure plates using the program Photoshop version 8.0	Photoshop		Suggestion: Adobe Photoshop (RRID:SCR_014199)(link)

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq