Document Identifier: **PMC8552857**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Materials and Methods Peripheral blood mononuclear cell culture The experiment procedure was approved by the Ethics Committee of Faculty of Medicine of University of Sao Paulo (FMUSPCAAE approval No. 52622616.8.0000.0065) and conformed to the guidelines of the Declaration of Helsinki.
Consent: All participants filled in the informed consent.
Euthanasia Agents: PBMC culture maintenance , hydrogen peroxide and antipsychotic treatment The PBMCs were maintained in a 6-well dish , in RPMI 1640 medium supplemented with 10 bovine fetal serum (Sigma) and antibiotics (penicillinstreptomycin 1 vv) at 37degC and an atmosphere with 5 of CO2 .
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.

Replication
For each sample , a 100 mL aliquot was transferred to a 96-well dish , in triplicate and added 10 mL of MTT reagent was added.
Number: For each sample , a 100 mL aliquot was transferred to a 96-well dish , in triplicate and added 10 mL of MTT reagent was added.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The diluted blood was carefully transferred to a conic tube containing 6 mL of Ficoll (Sigma, Sigma, St. Louis, MO, USA).	Sigma		
Antipsychotic preparation The antipsychotics were diluted in the following way: haloperidol (Sigma): 2.5 mg in 1.2 mL of DMSO (Sigma) clozapine (Biogen , Cambridge , MA , USA): 25 mg in 1 mL DMSO aripiprazole (Sigma): 2.5 mg in 500 mL PBS.	Biogen		Suggestion: Biogen Idec (RRID:SCR_003790)(link)
	Sigma)		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Differences in TL between control and vehicle within each treatment were investigated with one-way analysis of variance (ANOVA) followed by Tukeys post hoc test.	analysis of variance
	ANOVA
	Tukeys post hoc test
Oligonucleotides	
The primer sequences (5-3) were: telomere forward: GGT TTT TGA GGG TGA GGG TGA GGG TGA GGG TGA GGG T telomere reverse: TCC CGA CTA TCC CTA TCC CTA TCC CTA TCC CTA TCC CTA 36B4 forwardreverse CAG CAA GTG GGA AGG TGT AAT CC 36B4 forwardreverse CCC ATT CTA TCA TCA ACG GGT ACA A .	GGT TTT TGA GGG TGA GGG TGA GGG TGA GGG TGA GGG T telomere
	TCC CGA CTA TCC CTA TCC CTA TCC CTA TCC CTA TCC CTA 36B4 forwardreverse CAG CAA GTG GGA AGG TGT AAT CC 36B4
	forwardreverse
	CCC ATT CTA TCA TCA ACG GGT ACA A

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq