Document Identifier: **PMC8556563**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: This study followed all the guidelines for experimental investigation in human subjects and was approved by the Samsung Medical Center Institutional Review Board and adhered to the tenets of the Declaration of Helsinki.
Inclusion and Exclusion Criteria
The inclusion criteria of OAG included (1) the presence of glaucomatous optic disc changes, such as increased cupping, diffuse or focal neural rim thinning, disc hemorrhage, or RNFL defect; (2) an open angle on gonioscopic examination; and (3) glaucomatous visual field (VF) defects detected by more than one reliable test for at least two of the following three criteria: (1) a cluster of three points with a probability of less than 5% on the pattern deviation map in at least one hemifield, including at least one point with a probability of less than 1% or a cluster of two points with a probability of less than 1%; (2) a glaucoma hemifield test result outside normal limits; and (3) a pattern standard deviation (PSD) of 95% outside the normal limits.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: The inclusion criteria of NAION included (1) development of painless vision loss within 30 days of documented optic disc swelling confirmed by an ophthalmologist, (2) patient age between 40 and 85 years, (3) VF loss consistent with NAION, and (4) a relative afferent pupillary defect, unless the fellow eye affected in the past or both eyes were simultaneously affected with similar intensity.
Randomization
One eye per patient was used, that is, the randomly selected eye in the OAG group and the affected eye in the NAION group.
Blinding
Two independent readers (EJL and HJK) who were blinded to the clinical information of the patients evaluated the presence of MvD, and discrepancies were resolved by consensus.
Power Analysis

not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The area covered by the large retinal vessels was included as part of the MvD if the dropout extended beyond the vessels. The en face OCT image with marked MvD was then manually superimposed on the fundus photograph using a commercial software (Photoshop CC; Adobe Systems, Mountain View, CA, USA; Fig. 1).	Photoshop		Suggestion: Adobe Photoshop (RRID:SCR_014199)(link)
Measurements were performed using ImageJ (National Institutes of Health, Bethesda, MD, USA).	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
All statistical analyses were performed using SPSS statistical software (version 25.0; SPSS, Chicago, IL, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Eyes with media opacities and systemic or ocular diseases that could affect the VF test and OCTA were excluded.	VF test
Quantitatively, we correlated the width/ location of MvD, RNFL defect, βPPA, and βPPA–RNFL defect superimposition using Spearman's correlation analysis.	Spearman's correlation
18 Categorical variables were compared using Fisher's exact test.	Fisher's exact test
Regarding continuous variables, we used an independent t-test to compare the OAG and NAION groups, and one-way ANOVA and ad hoc analysis using Tukey's method were used to compare across the subgroups.	t-test
	ANOVA
	Tukey's method

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq