Document Identifier: **PMC8564844**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: We received approval from the Ethics Committee of the Eye Department, Clinical Centre of Vojvodina and the Teaching-Scientific Council of the Faculty of Medicine in Novi Sad.
Consent: A written plan with the informed consent form and patient information was approved by the ethics commissions of the Faculty of Medicine in Novi Sad.
Inclusion and Exclusion Criteria
Inclusion criterion was diagnosis of primary open-angle glaucoma on maximal antiglaucoma drug therapy with uncontrolled eye pressure.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
Patients and Methods This prospective, randomized, controlled clinical study included 40 patients (40 eyes) who had unregulated eye pressure despite taking maximal antiglaucoma pharmacotherapy, i.e., patients with refractory glaucoma, treated at the Eye Department, Clinical Center of Vojvodina, Novi Sad, Serbia, between January 2016 and June 2019.
Blinding
not detected.
Power Analysis
The following diode infrared laser (810 nm wavelength) parameters were used: power of 2.5 W and duration of 1.5 seconds.

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Statistical analysis was performed using the SPSS version 23.0 software.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Methods of identification of empirical distributions, methods for estimating equality of variance (Levenes test) and methods for estimating significance of difference were used.	Levenes test
T test was used for repeated measurements if assumption of the equality of variance was fulfilled, t-test for independent samples, and if not, then Welchs test was used.	T test
	t-test
	Welchs test
Also, Friedman test was used for comparisons within one study group, whereas Mann-Whitney test was used for comparisons between the study groups.	Friedman test
	Mann-Whitney test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq