Document Identifier: **PMC8568023**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The study was approved by the Bioethics Committee, approval no. KB1472017.
Inclusion and Exclusion Criteria
Exclusion criteria included other severe chronic diseases or delayed psychomotor development, IgAN in the course of systemic diseases, and IgAVN without kidney biopsy (these patients are not included in the registry), as well as age over 18 years (by December 30, 2018) and follow-up shorter than 3 months (for patients in the registry).
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: All children who were under 18 years of age by December 30, 2018, with at least a 3-month follow-up after kidney biopsy with urine and serum samples available, were included in the study.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Lilliefors and Shapiro-Wilk tests were used to test the normality of the data.	Shapiro-Wilk tests
Next, parametric tests for normally distributed data were used: Students t-test for independent samples (comparison of 2 groups) and ANOVA analysis of variance for the comparison of 3 groups.	t-test
	ANOVA
Non-parametric tests were used for non-normally distributed variables: Mann-Whitney (2 groups) and Kruskal-Wallis (3 groups).	Mann-Whitney
	Kruskal-Wallis
The statistical significance of the increment of individual variables was evaluated using the Students t-test for dependent samples and the Wilcoxon test (normal and non-normal distributions, respectively).	Wilcoxon test
To compare IgAVN vs. IgAN for the variables urine TNFR, serum TNFR, and log serum Gd-IgA1 between IgAVN and IgAN groups and correlate these variables with age at onset, height, and weight, ANCOVA covariance analysis was performed.	ANCOVA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq