Document Identifier: **PMC8572138**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The procedures for this study were conformed to the Helsinki protocol for clinical trials and the study was approved by the local regional Ethics Committee (ANBU 032018).
Inclusion and Exclusion Criteria
BCSs were enrolled if they met the following inclusion criteria: <= 5 years after surgery for BC (mastectomy andor quadrantectomy with or without axillary surgery) age from 30 to 70 years >= 6 months since completion chemotherapy not physically active (i.e., not meeting the physical activity guidelines for cancer survivors in the previous 6 months: 150 minweek of moderate intensity aerobic exercise and two daysweek of muscle-strengthening exercise 14).
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: BCSs were enrolled if they met the following inclusion criteria: <= 5 years after surgery for BC (mastectomy andor quadrantectomy with or without axillary surgery) age from 30 to 70 years >= 6 months since completion chemotherapy not physically active (i.e., not meeting the physical activity guidelines for cancer survivors in the previous 6 months: 150 minweek of moderate intensity aerobic exercise and two daysweek of muscle-strengthening exercise 14).
Randomization
not detected.
Blinding
not detected.
Power Analysis
2.7 Sample size calculation and data analysis The sample size was calculated using the Wilsons confidence interval method 58 with 95 confidence level, which resulted in 10 subjects per group who were needed to evaluate an adherence of 90 (i.e., 9 over 10 subjects who maintain, at t3, the PA level recommended by the

exercise professional and based on current exercise guidelines for secondary prevention 13, 14 with an interval width of 0.9 (95 CI: 0.59-0.98).
Replication
The circuit was repeated 2 times and each round always started with the aerobic exercise.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Within-group changes were checked using the Wilcoxon rank test between each time point.	Wilcoxon rank test
Intergroup differences were assessed using the Mann-Whitney U test to compare data between the two groups at each time point.	Mann-Whitney U test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq