Document Identifier: **PMC8577671**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Methods Ethical approval This study was approved by Medical Ethics Committee of the First Affiliated Hospital of Anhui Medical University (No. 2019-022).
Consent: Written informed consents were signed by all participants.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Clinical specimens Thirty-five specimens of lung adenocarcinoma tissues were collected from patients (21 males and 14 females, aged from 22 to 68 years, with an average age of 44.1 +- 12.0 years) who were pathologically diagnosed with lung adenocarcinoma and underwent resection of lung adenocarcinoma tumors in the First Affiliated Hospital of Anhui Medical University between April 2016 and February 2017.
Subject Demographics
Age: Clinical specimens Thirty-five specimens of lung adenocarcinoma tissues were collected from patients (21 males and 14 females, aged from 22 to 68 years, with an average age of 44.1 +- 12.0 years) who were pathologically diagnosed with lung adenocarcinoma and underwent resection of lung adenocarcinoma tumors in the First Affiliated Hospital of Anhui Medical University between April 2016 and February 2017.
Randomization
Finally , five fields were randomly selected and ImageJ (National Institutes of Health , Bethesda , MD , USA) was used to calculate the number of colonies .
Blinding
not detected.
Power Analysis
Sample size was determined using G power for power analysis.

Replication
Number: Data are exhibited as the mean +- standard deviation from at least three independent assays.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The GAPDH antibody was utilized for normalization .	GAPDH		
The primary antibodies are exhibited as follows: RhoA (Abcam , Cambridge , UK , ab54835) rho associated coiled-coil containing protein kinase 1 (ROCK1 , Abcam , ab45171	ROCK1		
Experimental Models: Cell Lines			
Cell culture and treatment Cell Bank of the Chinese Academy of Sciences (Shanghai, China) provided the human lung adenocarcinoma cell lines (A549, H1650, PC-9, and H441) and a normal human bronchial epithelial cell line (BEAS-2B).	A549		Suggestion: RRID:CVCL_0023(link)
	PC-9		Suggestion: RRID:CVCL_B260(link)
Cotransfection of RhoA 3-UTR WT vector or Mut vector and miR-345-5p mimics or NC mimics into PC-9 and H1650 cells was performed using Lipofectamine 2000 transfection reagent (Invitrogen) .	H1650		Suggestion: RRID:CVCL_1483(link)
Recombinant DNA			
GCUGACUCCUAGUCCAGGGCUC) , NC inhibitor , empty pcDNA3.1 , and pcDNA3.1ras homolog family member A (RhoA) (RiboBio , Guangzhou , China) were transfected into lung adenocarcinoma cells for 48 h as per the product manuals .	pcDNA3.1		Suggestion: RRID:Addgene_100544(link)
	pcDNA3.1ras		
Mutant (Mut) and wild-type (WT) sequences of RhoA 3-UTR were amplified and subcloned into the pmirGLO luciferase reporter vectors (Promega , Madison , WI , USA)	pmirGLO		
Software and Algorithms			
Sample size was determined using G power for power analysis.	G power		
Finally , five fields were randomly selected and ImageJ (National Institutes of Health , Bethesda , MD , USA) was used to calculate the number of colonies .	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Bioinformatics analysis Based on the PITA (http:genie.weizmann.ac.ilpubsmir07mir07_data . html) and PicTar (https:pictar .	PITA		Suggestion: PITA (RRID:SCR_010853)(link)
	PicTar		

mdc-berlin . de) online databases which predict potential targets of miR-345-5p , 23 mRNAs were identified.			
Statistical analysis We used the SPSS 20.0 (St Louis, MO, USA) for the statistical analysis.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Levene test was used for homogeneity of variance test, and Shapiro-Wilk test was used for normal distribution test.	Levene test
	Shapiro-Wilk test
Comparison between two groups was conducted with Students t test, and one-way analysis of variance followed by Dunnett post hoc test or Tukey post hoc test was adopted for comparison among multiple groups.	Students t test
	analysis of variance
	Dunnett post hoc test
	Tukey post hoc test
Spearman correlation analysis was applied to analyze the expression correlation between miR-345-5p and RhoA in lung adenocarcinoma tissues.	Spearman correlation

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq