Document Identifier: **PMC8580088**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: 2 MATERIALS AND METHODS 2.1 ACVR1 screening Peripheral blood samples were collected upon informed consent administration.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Samples were measured in duplicate and the results were normalized on the geometric mean of B2M and HPRT genes.
Number: Samples were measured in duplicate and the results were normalized on the geometric mean of B2M and HPRT genes.

Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Primary antibodies were diluted as follows: P-SMAD159 1:2000 (13820, SMAD1 1:2000 9743, Cell Signaling) anti-FLAG M2 monoclonal antibody 1:3000 (F1804, Sigma-Aldrich, Merk) anti-GAPDH 1:20000 (MAB374, Millipore).	SMAD1		
	anti-FLAG		
	anti-GAPDH		
Experimental Models: Cell Lines			
ID1 gene expression and P-SMAD159 proteins levels were evaluated by seeding 3 x 105 U2OS in 6-well plates for transfection with 2 ug of pCMV_ACVR1 constructs using the Lipofectamine 2000 reagent (Invitrogen, Life Technologies).	U2OS		Suggestion: RRID:CVCL_0042(link)
Recombinant DNA			
2.2 In silico analysis In order to estimate the impact of the identified variant on ACVR1 receptor function, we performed an in silico analysis by using the Ensembl Variant Effect Predictor (VEP) toolset (available at https:grch37.ensembl.orgToolsVEP) McLaren et al., 2016. 2.3 Expression plasmids Complementary DNA (cDNA) for human, wild-type ACVR1 (Table S1 for oligonucleotides used to generate the product) was obtained by RT-PCR and then subcloned into the expression vector pCMV-3Tag-8 (pCMV_ACVR1) without the native STOP codon, to be in frame with the 3xFLAG tag provided by the vector.	pCMV-3Tag-8		
The generation of the pGL4.17 vector containing the BMP-Responsive Element (BRE) upstream the Luciferase reporter gene (indicated as pGL4.17-BRE) was previously described by our group (Cappato et al., 2016).	pGL4.17		
	pGL4.17-BRE		Suggestion: RRID:Addgene_45126(link)
Transient transfections were performed in 96-well plates, by seeding 5 x 104well U2OS cells directly in the transfection mix, composed by 30 ng of the pGL4.17-BRE, 40 ng of pCMV_ACVR1 constructs and 2.5 ng of pGL4.73hRlucSV40	pGL4.73hRlucSV40		

to normalize toward transfection efficiency, using the Lipofectamine 2000 reagent protocol (Invitrogen, Life Technologies).			
ID1 gene expression and P-SMAD159 proteins levels were evaluated by seeding 3 x 105 U2OS in 6-well plates for transfection with 2 ug of pCMV_ACVR1 constructs using the Lipofectamine 2000 reagent (Invitrogen, Life Technologies).	pCMV_ACVR1		
Software and Algorithms			
Genomic DNA fragments corresponding to the nine coding exons and flanking intronic sequences of ACVR1 (RefSeq NM_001111067.4 RefSeq NP_001104537.1), were amplified by polymerase chain reaction (PCR the primer sequences are listed in Table S1) using the GoTaq Master mix (Promega) with reactions carried out in a GeneAmp PCR System 2720 Thermocycler (Applied Biosystems).	RefSeq		
Sequence variations were described according to the Human Genome Variation Society guidelines (http:www.hgvs.org).	http:www.hgvs.org		
2.2 In silico analysis In order to estimate the impact of the identified variant on ACVR1 receptor function, we performed an in silico analysis by using the Ensembl Variant Effect Predictor (VEP) toolset (available at https:grch37.ensembl.orgToolsVEP) McLaren et al., 2016. 2.3 Expression plasmids Complementary DNA (cDNA) for human, wild-type ACVR1 (Table S1 for oligonucleotides used to generate the product) was obtained by RT-PCR and then subcloned into the expression vector pCMV-3Tag-8 (pCMV_ACVR1) without the native STOP codon, to be in frame with the 3xFLAG tag provided by the vector.	Ensembl Variant Effect Predictor		
	Variant		Suggestion: VARIANT (RRID:SCR_005194)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The Unpair t test was applied to verify statistical significance of the observed variations (https:www.graphpad.comquickcalcsttest1.cfm).	t test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq