Document Identifier: **PMC8593692**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: An informed consent form was filled by all participants after explaining the aims of the study.
IRB: Moreover, the Ethics Committee of Tehran University of Medical Sciences approved the protocol of this study.
Inclusion and Exclusion Criteria
We excluded patients with recent (through six months) heart failure, myocardial infarction, or stroke in addition to participants with chronic conditions or rheumatologic diseases, and pregnant women.
Attrition
not detected.
Sex as a biological variable
Study Participants A total of 5115 obeseoverweight adults involving 4042 (79.1) females and 1073 (20.9) males referred to the weight loss clinic from June 2017 to June 2019 of whom 1870 were overweight and the remaining 3245 were obese.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication

not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Hypertension Definition The diagnosis of hypertension was made based on BP > 13080 on two separate office visits or previous treatment with antihypertensive drugs according to the American College of Cardiology (ACC) and the American Heart Association (AHA) guideline for the Prevention, Detection, Evaluation and Management of High Blood Pressure in Adults.	American Heart Association		Suggestion: American Heart Association (RRID:SCR_007210)(link)
35 All statistical analyses were performed using IBM SPSS Statistics version 20.0	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical Analysis The independent t-test and chi-square test were employed to compare quantitative and categorical variables , respectively between normotensive and hypertensive patients .	t-test
	chi-square test
Pearsons correlation test was used to examine the association of systolic blood pressure and anthropometric indices .	Pearsons correlation test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq