Document Identifier: **PMC8594505**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: Asked to remain usual PA levels and received the intervention upon completion of the study 1) Quality of life (SF-36 and IBCSG-QLC) and 2) fatigue (FACT-F) FACT-F Pinto et al. (33) n = 46 (I:20 , C: 26) 57.3 (9.7) 56 female , 98 Caucasian , 76 higher education Colon cancer (57) and rectal cancer (43) Surgery 100 Chemo 83 Radiation 43 3.0 (1.6) yr a Inclusion: 1) age >=18 yr , 2) completed primary and adjuvant treatments for colon or rectal cancer , 3) <=5 yr since treatment completion , 4) able to read and speak English , 5) consent for medical chart review , 6) able to walk unassisted , 7) sedentary (exercising <60 min*wk-1 at moderate intensity or <20 min*wk-1 of vigorous intensity over the past 6 months) , 8) access to a telephone Exclusion: 1) history of cancer or 2) a medical or current psychiatric illness
Inclusion and Exclusion Criteria
Eligibility criteria Randomized controlled trials meeting the following criteria were included: (i) study sample of cancer survivors (except for nonmelanoma skin cancer) (ii) age 18 yr or older (iii) completed cancer treatment (chemotherapy , surgery , and or radiotherapy being on adjuvant hormonal therapy was allowed) (iv) the intervention was designed to increase PA (v) the intervention did not target other health behaviors (e.g. , smoking or nutrition) (vi) the intervention was home-based (vii) PA was of low to moderate intensity (viii) PA was not personally supervised and did not include a structured training regime , such as anaerobic training or yoga (ix) the control group received usual care or a nonphysical intervention (x) symptoms of fatigue , depression , or anxiety were assessed with a validated questionnaire , a subscale from a validated questionnaire , or a visual analog scale (VAS) and (xi) scores for the questionnaires were reported.
Attrition
not detected.
Sex as a biological variable
Study Sample Size Age (yr) , Female , Caucasian , Higher Education Diagnosis ( )
Subject Demographics
Age: Eligibility criteria Randomized controlled trials meeting the following criteria were included: (i) study sample of cancer survivors (except for nonmelanoma skin cancer) (ii) age 18 yr or older (iii) completed cancer treatment (chemotherapy , surgery , and or radiotherapy being on adjuvant hormonal therapy was allowed) (iv) the intervention was designed to increase PA (v) the intervention did not target other health behaviors (e.g. , smoking or nutrition) (vi) the intervention was home-based (vii) PA was of low to moderate intensity (viii) PA was not personally supervised and did not include a structured training regime , such as anaerobic training or yoga (ix) the control group received usual care or a nonphysical intervention (x) symptoms of fatigue , depression , or anxiety were assessed with a validated questionnaire , a subscale

from a validated questionnaire , or a visual analog scale (VAS) and (xi) scores for the questionnaires were reported.
Randomization
A combination of MeSH terms and free-text words was used , including randomized controlled trial , cancer , physical activity , home-based or community-based , and fatigue , depression , or anxiety .
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Search strategy On May 19 , 2020 , we searched PubMed , CINAHL , PsycINFO , and Web of Science for relevant studies .	PubMed		Suggestion: PubMed (RRID:SCR_004846)(link)
	PsycINFO		Suggestion: PsycINFO (RRID:SCR_014799)(link)
A combination of MeSH terms and free-text words was used , including randomized controlled trial , cancer , physical activity , home-based or community-based , and fatigue , depression , or anxiety .	MeSH		Suggestion: MeSH (RRID:SCR_004750)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Heterogeneity in effect measures between studies was assessed using the I 2 statistic.	I 2 statistic

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq