Document Identifier: **PMC8602394**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not detected.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
After randomization, normalization, and FCS file generation, the files were further analyzed in Cytobank (Beckman Coulter, Brea, CA).
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The blotted proteins were reacted to the indicated antibody (anti-3A5 in Figure S1 , and anti-H18 in Figure S2) with rotation at 21 degC for 3 h .	anti-H18		
Washing was performed with PBST three times for 10 min , followed by incubation with HRPO-conjugated anti-mouse antibody (Catalog No. 62-6520 , ThermoFisher Scientific) .	anti-mouse		
The negative control monoclonal antibody was a mouse IgG1 (clone T248	mouse IgG1		
The secondary antibody , Alexa Fluor 488-labeled anti-mouse IgG (Catalog No. AP124JA4 , Sigma-Aldrich) was added (1:1000) .	Alexa Fluor 488-labeled anti-mouse IgG		
	anti-mouse IgG		
anti-GFP (Catalog No. A11122 , ThermoFisher Scientific 43) , and anti-CD45 (Catalog No. 3089003B , Fluidigm , South San Francisco , CA 44) , and then conjugated in house using Maxpar Antibody Labeling Kit (Fluidigm) according to the instructions of the manufacturer .	anti-CD45		
The following antibody concentrations were used: anti-H2: 5 ugml , anti-H3: 5 ugml , anti-H18: 5 ugml , anti-L1: 1 ugml , anti-L2: 7.5 ugml , anti-L4: 7.5 ugml , anti-L6: 10 ugml , anti-IgM: 10 ugml , anti-P1: 7.5 ugml , anti-3A5: 5 ugml , and anti-GFP: 10 ugml .	anti-H2		
	anti-H3		
	anti-L1		
	anti-L2		
	anti-L4		
	anti-L6		
	anti-P1		
	anti-3A5		
	anti-GFP		
The L6 (IgM isotype) marker was measured indirectly via metal tag-labeled anti-IgM antibody .	L6 (IgM		
The negative control monoclonal antibody was a mouse IgG1 (clone HI30 , anti-human 89 Y-labeled CD45) in 1:100 dilution .	anti-human 89 Y-labeled CD45		
	Y-labeled		
The isotypes of anti-Drosophila antibodies were determined by the IsoStrip(tm) Antibody Isotyping Kit (Catalog No. 11493027001 , Roche ,	anti-Drosophila		

Basel , Switzerland) according to the instructions of the manufacturer .			
Cells were washed with 200 ul MCSB , stained with anti-IgM antibody (volumes were the same as in the surface staining) , and incubated on ice for 30 min .	anti-IgM		
Experimental Models: Organisms/Strains			
Materials and methods Drosophila stocks The following Drosophila lines were used in the current study: w1118 (BSC9505, Bloomington Drosophila Stock Center, Bloomington,	w1118		Suggestion: RRID:BDSC_3605(link)
IN) , Ore-R (wild type) , w hdcD84TM3 , Kr>GFP 30 , lz-Gal4 , UAS-GFP + + (a gift from Bruno Lemaitre , Lausanne , Switzerland) 32	Ore-R		
	w hdcD84TM3 , Kr>GFP 30 , lz-Gal4 , UAS-GFP + +		
Software and Algorithms			
The microscopic analysis was carried out using a Zeiss Axioskope 2MOT epifluorescent microscope and Axiovision 2.4 software (Zeiss , Oberkochen , Germany) .	Axiovision		
After 45 min incubation at 4 degC , the cells were washed three times with ice-cold Schneiders medium and acquired on FACSCalibur (Beckton Dickinson , Franklin Lakes , NJ ,)	FACSCalibur		
After randomization, normalization, and FCS file generation, the files were further analyzed in Cytobank (Beckman Coulter, Brea, CA).	Cytobank		

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq