Document Identifier: **PMC8605897**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: This study was approved by the ethics committees of the Second Affiliated Hospital, Jiaxing University (Ethics number: jxey-2018038), which accorded with the ethical standards formulated in the Helsinki Declaration.
IACUC: The experiments were approved by the Animal Ethics Committee of Jiaxing University, Jiaxing, China. Statistical Analysis All experimental data were expressed as the mean ± standard deviation (S.D.).
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Materials and Methods Patients and Samples A total of 47 colon cancer patients (27 males and 20 females, aged from 45 to 81 years old, median age 63 years old) were recruited in this study who hospitalized in the Second Affiliated Hospital of Jiaxing University between 2018 Jul to 2020 Dec.
Subject Demographics
Age: Materials and Methods Patients and Samples A total of 47 colon cancer patients (27 males and 20 females, aged from 45 to 81 years old, median age 63 years old) were recruited in this study who hospitalized in the Second Affiliated Hospital of Jiaxing University between 2018 Jul to 2020 Dec.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication

not required.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Table 1 Demographic, Sex, and Clinical Stages of Patients with Colon Cancer and Healthy Donors Healthy Donors Colon Cancer Patients Total samples 20 47 Age (years) Min 42 45 Max 75 81 Mean 55.60 ± 2.24 62.62 ± 1.46 Sex Male 12 27 Female 8 20 Clinical stages Stage I–II – 16 Stage III–IV – 31 Flow Cytometry The human fluorescent-labeled antibodies included PerCP-Cy5.5-anti-CD3, FITC-anti-CD4, APC-Cy7-anti-CD8, APC-anti-CD161 and PE-anti-TCR Vα7.2.	APC-Cy7-anti-CD8		
	APC-anti-CD161		
The mouse fluorescent-labeled antibodies included FITC-anti-CD3, APC-Cy7-anti-CD8, APC-anti-CD24 and PE-anti-TCRβ.	PE-anti-TCRβ		
Then the slides were incubated with 1:200 diluted anti-CD161 antibody (ab197979, Abcam, USA) at 4 °C for overnight.	anti-CD161		Suggestion: (DSHB Cat# anti-CD38 (HB-7), RRID:AB_3105832)(link)
Slides were treated with secondary antibody anti-rabbit IgG (Servicebio, China) for 30 min and then incubated with simple stain DAB solution (Servicebio, China).	anti-rabbit IgG		
Experimental Models: Cell Lines			
BALB/c mice were inoculated subcutaneously in a single-cell suspension with 2.5×105 CT26 tumor cells.	CT26		Suggestion: RRID:CVCL_7254(link)
Experimental Models: Organisms/Strains			
Animal Experiments 8-week-old female BALB/c mice were purchased from Shanghai Laboratory Animal Center, Chinese Academy of Sciences (SLAC, CAS).	BALB/c		Suggestion: RRID:MGI:2683685(link)
Software and Algorithms			
The data were analyzed by FlowJo v7.6.2 software (Tree Star, Ashland, OR).	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)
TCGA and GEPIA Database Analysis The gene expressions of KLRB1 in tumor tissues, normal tissues, as well as different TNM stages of tumor	https:// cancergenome.nih	gov/	

tissues were studied by analyzing colon cancers of the TCGA database (https://cancergenome.nih.gov/).			
Statistical analyses were performed using GraphPad Prism 6 (GraphPad Software, USA).	GraphPad Prism		Suggestion: GraphPad Prism (RRID:SCR_002798)(link)
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical significance of expression changes were calculated by unpaired t-test for two groups.	t-test
The two-way ANOVA was employed to compare grouped and paired data sets as appropriate.	ANOVA
Correlations were analyzed using the Spearman’s rank correlation test.	Spearman’s rank correlation test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq