Document Identifier: **PMC8612137**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: MATERIALS AND METHODS The study was approved by the Joint Chinese University of Hong Kong-New Territories East Cluster Clinical Research Ethics Committee (reference number CREC-2015.327).
Consent: Informed consent was obtained 5 days before antibiotic completion, according to the ISPD guideline, and patients were then randomized to receive 1 additional week of the effective antibiotic treatment on top of the ISPD guideline (the extended group) or completion of antibiotics according to the guideline with no additional treatment (the standard group).
Inclusion and Exclusion Criteria
Patients with fungal peritonitis or obvious surgical problems that required laparotomy were excluded.
Attrition
The original plan was to recruit 360 patients who fulfilled the diagnostic criteria of PD peritonitis, which was based on at least two of the following 10: abdominal pain or cloudy PD effluent, leukocytosis in PD effluent (white cell count >100mL) and positive Gram stain or culture from PD effluent.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
Case selection This was a prospective randomized controlled study.
Blinding
not detected.
Power Analysis
A sample size of 300 would have 80 power to detect such a difference.

Replication	
not required.	
Protocol Information	
Identifiers: The study was registered at ClinicalTrials.gov (NCT02593201).	NCT02593201

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Statistical analyses Statistical analyses were performed using SPSS Statistics version 24.0 (IBM, Armonk, NY, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Baseline demographic and clinical data were compared between the extended and standard groups by Students t-test, Mann-Whitney U-test or chi-square test as appropriate.	t-test
	Mann-Whitney U-test
	chi-square test
We performed exploratory analysis on PD effluent bacterial DNA levels before and after treatment and the data were compared by Wilcoxon signed-rank test.	Wilcoxon signed-rank test
Data were further analyzed by the intention-to-treat approach with Kaplan-Meier plot and univariate Cox regression analysis, with patient death, fungal or tuberculous peritonitis or catheter removal treated as censoring events.	Cox regression

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq