Document Identifier: **PMC8627652**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not required.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
MS survey scans were performed for mass-to-charge ratios ranging from 350-1,800, and the 20 most intense ions from the survey scans were analyzed by MSMS spectra in the LTQ, which were determined using Xcalibur mass spectrometer software in real-time.	MSMS		Suggestion: MSMS (RRID:SCR_003532)(link)
Processing of MS data Raw files were processed using MaxQuant (version 1.2.2.5) (Cox & Mann, 2008) and searched against the UniProt Staphylococcus NCTC8325 protein database (20151213 UniProtKB: UP000008816 2,889 sequences).	MaxQuant		Suggestion: MaxQuant (RRID:SCR_014485)(link)
	UniProt		Suggestion: UniProt (RRID:SCR_002380)(link)
Bioinformatics analysis of the endogenous peptidome To annotate the features of the peptidome, EMBOSS (Rice, Longden & Bleasby, 2000) was used to calculate the isoelectric point (pI) and molecular weight (Mw) of the peptide sequences.	EMBOSS		Suggestion: EMBOSS (RRID:SCR_008493)(link)
To analyze the sequence patterns of proteases to generate endogenous peptides, the amino acid sequences surrounding the termini of peptides were extracted, and overrepresented amino acids were analyzed using iceLogo (Colaert et al., 2009).	iceLogo		Suggestion: iceLogo (RRID:SCR_012137)(link)
Interactions among precursor proteins were annotated by the STRING database (Szklarczyk et al., 2017), and the protein-protein interaction network was constructed using Cytoscape software (Kohl, Wiese & Warscheid, 2011).	STRING		Suggestion: STRING (RRID:SCR_005223)(link)
	Cytoscape		
The database for annotation, visualization, and integrated discovery (DAVID) (version 6.8) (Huang da, Sherman & Lempicki, 2009) was used to identify significantly enriched gene ontology terms in biological processes (BP), molecular function (MF), and cellular component (CC) categories for the precursor proteins of the DE endogenous peptidome.	DAVID		Suggestion: DAVID (RRID:SCR_001881)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
One-way ANOVA was performed to analyze the differential expression of peptides among groups, and a peptide was considered differentially expressed if the FDR-q value was less than 0.01.	ANOVA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq