Document Identifier: **PMC8631182**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Materials and Methods Study Design and Participants This observational cross-sectional study was approved by the King Abdulaziz University (KAU)-Research Ethics Committee (approval number: 609-18) and was performed in accordance with the Helsinki Declaration.
Consent: Written informed consents were taken from all participants.
Inclusion and Exclusion Criteria
The exclusion of subjects from the study was built on the history, clinical picture, and investigations.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
The chronic hepatitis patients were randomly recruited from Gastroenterology and Hepatology Department, King Abdulaziz University (KAU) Hospital between January 2018 and December 2020.
Blinding
not detected.
Power Analysis
To calculate a cost-effective sample size, we set the significance level at $P < 0.05$ and used a statistical power of 80 which is widely used to avoid false-negative associations.
Replication

not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Then, streptavidin-HRP is added to bind to the biotinylated NT antibody.	NT		
Software and Algorithms			
The HV serology markers were measured by the chemiluminescent microparticle immunoassay method (Abbott Diagnostics, IL, USA) and serum levels of HBV DNA and HCV RNA were measured by the real-time PCR kits (Roche Diagnostics, IN, USA) according to the manufacturers protocols.	Abbott		Suggestion: Abbott (RRID:SCR_010477)(link)
16 Statistical Analysis The SPSS version 22 (SPSS Inc., IL, USA) was used.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The ANOVA with Tukeys test was used for multiple comparisons.	ANOVA
	Tukeys test
Pearson correlation was used to detect correlations between viral loads and HBsAG and glutamine or nitrotyrosine levels of hepatitis patients.	Pearson correlation
Moreover, Spearman correlation was used to detect correlations between glutamine and nitrotyrosine levels of normal subjects or hepatitis patients and the standard diagnostic tests for hepatitis.	Spearman correlation

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq