Document Identifier: **PMC8636821**

# SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

**Table 1: Rigor Adherence Table**

| |
|---|
| Ethics |
| Field Sample Permit: 2.2 PCL scaffold fabrication PEOPCL blend preparation and annealing were carried out as described 17 with minor modifications . |
| IACUC: 2.6 In vivo osteochondral repair model and histoprocessing All experiments with animals were carried out under protocols approved by the University of Montreal Animal Care Committee ( Protocol 16-100 , Polytechnique Montreal Protocol ANI-1516-20 ) and according to ARRIVE guidelines 34 |
| Euthanasia Agents: At 1 day ( N = 1 ) or 6 weeks ( N = 6 ) post-operative rabbits were euthanized under anesthesia by pentobarbital intravenous injection . |
| Inclusion and Exclusion Criteria |
| not detected. |
| Attrition |
| not detected. |
| Sex as a biological variable |
| Skeletally mature retired breeder rabbits ( N = 7 , all females due to unavailable male retired breeders , over 10 months old , 4-5 kg , Charles River , St . Constant , QC , Canada ) were housed in individual cages . |
| Subject Demographics |
| Age: Skeletally mature retired breeder rabbits ( N = 7 , all females due to unavailable male retired breeders , over 10 months old , 4-5 kg , Charles River , St . Constant , QC , Canada ) were housed in individual cages . |
| Weight: Skeletally mature retired breeder rabbits ( N = 7 , all females due to unavailable male retired breeders , over 10 months old , 4-5 kg , Charles River , St . Constant , QC , Canada ) were housed in individual cages . |
| Randomization |
| Rabbits and knees were randomized prior to surgery to 4 conditions to obtain N = 6 osteochondral defects per condition . |
| Blinding |

| |
|---|
| 2.7 Micro-computed tomography Distal femurs were imaged with a Skyscan 1172 instrument ( 70 kV , 141 mA , pixel size 9 mm , aluminum filter 0.5 mm , frame averaging ( 3) , 0.42deg rotation step) . 3-D images were reconstructed using NRecon and analyzed by a blinded observer using DataViewer to align each drill hole perpendicular to the z-axis ( N = 28) . |
| Power Analysis |
| not detected. |
| Replication |
| not required. |

**Table 2: Key Resources Table**

| Your Sentences | REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|---|---|---|
| Software and Algorithms | | | |
| PSS , M w 70,000 , 30 wt in H2O , and PDADMAC , M w 100,000-200,000 , 20 wt in H2O ( 409014) , were purchased from Sigma ( Oakville , ON , Canada) . | Sigma | | Suggestion: sigma-tau (RRID:SCR_000488)( link) |
| An image analysis software ( SigmaScan Pro V5 , SigmaPlot ) was used to calculate the volume average pore size from 50 pore measures per sample. | SigmaPlot | | Suggestion: SigmaPlot (RRID:SCR_003210)( link) |
| Nuclei were counted in 10x magnification images of the blood clot at the edge of each scaffold and the middle of each clot ( ImageJ ) in 2 cryosections from 3 independent samples per condition . | ImageJ | | Suggestion: ImageJ, RRID:SCR_003070( link) |

**Other Entities Detected**

| Your Sentences | Recognized Entity |
|---|---|
| Statistical Tests | |
| 2-tailed Students t-test was used to analyze clot retraction , leukocyte density and transient knee effusion ( N = 3) . | Students t-test |
| Analysis of the variance of the mean ( ANOVA ) with Tukey Honest Significant Differences ( HSD ) post-hoc was used to analyze the effect of condition on osteochondral defect bleeding ( N = 7) , and the effect of scaffold on 6-week histological repair using microdrill hole as the experimental unit ( N = 6) . | ANOVA |
| | Tukey Honest |

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq