Document Identifier: **PMC8643178**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
Second, based on their clinical stages, UCEC samples can be divided into 375 patients with early stage (stage I and II) and 145 patients with advanced stage (stage III and IV), and then the relationship between expression and clinical stages was determined.
Sex as a biological variable
not required.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Each sample was acquired using the American Illumina HiSeq 2000 RNA sequencing platform, and the ENSEMBL system was used for gene annotation.	ENSEMBL		Suggestion: Ensembl (RRID:SCR_002344)(link)
Using the edgeR package in the R software, RNA-seq data of 552 UCEC samples and 35 normal endometrial samples were utilized to screen out differentially expressed genes (DEGs) with the cut-off criteria of log2FC>=1 and false discovery rate (FDR) <0.05.	edgeR		
Furthermore , DE miRNAs were combined to construct a ceRNA regulatory network by the following steps: (1) predicting the interaction between DE lncRNAs and DE miRNAs using miRcode (http:www.mircode.org) database (2) predicting the interaction between DE miRNAs and DE mRNAs using TargetScan (http:www.targetscan.org)	TargetScan		Suggestion: TargetScan (RRID:SCR_010845)(link)
), miRTarBase (http:miRTarBase.cuhk.edu.cn), and miRBase (http:www.mirbase.org) databases (3) selecting duplicate DE miRNAs as common connections.	http:www.mirbase.org		
Finally, we constructed ceRNA regulatory networks and visualized them using Cytoscape software (version 3.7.0).	Cytoscape		Suggestion: Cytoscape (RRID:SCR_003032)(link)
Construction of Protein-Protein Interaction (PPI) Network and Screening of Key Genes We uploaded DE mRNAs related to pathological tissue subtypes of UCEC identified in WGCNA to the Search Tool for the Retrieval of Interacting Genes (STRING, http:string-db.org) online analysis software to identify the potential relationship between proteins encoded by DE mRNAs.	STRING		
Molecular Complex Detection algorithm was performed using MCODE plug-in to filter out highly relevant modules from the entire PPI network and assigned each molecule to the MCODE value to calculate	MCODE		

clustering seed genes and clustered genes in the module.			
The CyTargetLinker plug-in based on Wikipathways database (https:www.wikipathways.orgindex.phpWikiPathways) was used to perform the functional pathway enrichment analysis of DE mRNAs in the constructed PPI network.	Wikipathways		Suggestion: WikiPathways (RRID:SCR_002134)(link)
In addition, to verify whether Wnt signaling pathway-related molecules predicted by the metabolic pathway analysis in PPI network had reliable biological significance, we used TCGA data in Oncomine (https:www.oncomine.org) database to observe the expression levels of genes related to Wnt signaling pathways in type I and type II UCEC, respectively.	Oncomine		
	https:www.oncomine.org		
The Human Protein Atlas (http:www.proteinatlas.org)23 was employed to map the tissue protein expression pattern for the key RNAs.	http:www.proteinatlas.org		
Statistical Methods The data were presented as mean +- standard deviation, and GraphPad Prism software was used for statistical analysis.	GraphPad Prism		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
An unsupervised co-expression relationship between genes was established using Pearson correlation coefficients .	Pearson correlation
Five hundred and twenty UCEC cases were divided into high- and low-expression groups to perform overall survival (OS) analysis by using the Log rank test method, determining the associations between key RNAs and OS rate.	Log rank test
The comparison between the two groups was performed by t test, and the comparison between multiple groups was performed by one-way analysis of variance (ANOVA).	t test
	analysis of variance
	ANOVA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq