Document Identifier: **PMC8646637**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: For control groups , blood samples were collected from severe ill ICU patients that presented an high sequential organ failure assessment (SOFA)-score (n = 9 , ICU control , Table S1 and Figure S1) , which predicts the risk of morbidity and mortality due to sepsis and COVID-19 as previously described , 12 as well as from healthy volunteers at the blood donation center in Tuebingen , after written consent was obtained. 2.2 Evaluation of the clinical data Clinical data related to treatments , demography and outcome were obtained from electronic medical records of the university hospital of Tuebingen .
IRB: The study protocol was approved by the Institutional Review Board of the University of Tuebingen.
Inclusion and Exclusion Criteria
2.6 Sera preparation and incubation with washed platelets All sera were heat-inactivated at 56degC for 30 min , followed by a sharp centrifugation step at 5000g to exclude any unspecific effects like the activation of platelets via complement and immune complexes .
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
For adhesion quantification , images were captured from five randomly chosen microscopic fields and analyzed with the CellSens Standard software (Olympus)
Blinding
not detected.
Power Analysis

not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Next , cells were stained with a monoclonal antibody (moAb) against CD62p (anti-CD62p 1:20 BD) and annexin-V (1:20 Immunotools) that detects externalized phosphatidylserine (PS) in the presence of 1 mM CaCl2 , (Sigma) for 15 min at RT in the dark .	CD62p		
	annexin-V		
2.7 IgG preparation and assessment of antibody-mediated signaling A commercially available IgG-purification-kit (MelonTM-Gel IgG Spin Purification Kit Thermo Fisher Scientific) was used for IgG purification as described in the manufacturers recommendations (for details , see Appendix) .	MelonTM-Gel IgG		
Finally , the membranes were incubated with rabbit phospho-AKT antibody (1:1000 Invitrogen) , mouse AKT antibody (1:1000 Abcam , Signaling) , rabbit pPI3K (p85 , N-SH , 1:1000 Merck) and mouse a-Tubulin (Cell Signaling) which was followed by the incubation with appropriate secondary anti-rabbit IRDye(r)800 and anti-mouse antibody IRDye(r)680 (1:3000 , LI.	phospho-AKT		
	mouse AKT		
	N-SH		
	mouse a-Tubulin		
	anti-rabbit		
	anti-mouse		
Recombinant DNA			
2.4 Platelet signaling pathway assay For the detection of AKT phosphorylation on Ser473 in platelets , whole blood samples were fixed using the commercially available PerFix-nc kit (Beckmann Coulter) and subsequently stained with CD41-PC5 (BD) , pAKT-PE and AKT-APC (both Invitrogen) at RT for 30 min .	pAKT-PE		
Software and Algorithms			
Relative intensity of western blots was quantified using ImageJ software (NIH) . 2.9 Microscopy and platelet adhesion test For the assessment of the adhesion ability of platelets , an adhesion assay was performed.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
For adhesion quantification , images were captured from five randomly	CellSens		

chosen microscopic fields and analyzed with the CellSens Standard software (Olympus)			
Statistical analysis was performed using GraphPad Prism, Version 8.1 (GraphPad).	GraphPad Prism		
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Non-parametric tests were used when data failed to follow a normal distribution as assessed by the DAgostino and Pearson omnibus normality test.	Pearson omnibus normality test
Group comparison was performed using the Wilcoxon rank-sum test.	Wilcoxon rank-sum test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq