Document Identifier: **PMC8647233**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: This original study for the EGFR-TKI resistance in lung cancer was approved by the Peking University Cancer Hospital Ethics Committee , and written informed consent was obtained from all participants prior to sample collection .
Consent: This original study for the EGFR-TKI resistance in lung cancer was approved by the Peking University Cancer Hospital Ethics Committee , and written informed consent was obtained from all participants prior to sample collection .
Inclusion and Exclusion Criteria
not detected.
Attrition
Five databases were included in the analysis , including PicTar , DIANA , TargetScan , miRanda and miRDB , wherein miRNA-mRNA interactions were reported consistently in at least two of the five databases .
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
Afterwards , the enriched circRNAs were amplified and transcribed into fluorescent cDNA using a random priming method (Arraystar Super RNA Labeling Kit , Arraystar) .
Blinding
not detected.
Power Analysis
not detected.

Replication
2.9 Statistical Analysis All experiments were performed in triplicates, wherein data are presented as mean +- standard deviation.
Number: 2.9 Statistical Analysis All experiments were performed in triplicates, wherein data are presented as mean +- standard deviation.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
2.2 Cell Lines and Cell Culture PCER cells were cultivated from parental PC9 cells after 6 months of gefitinib exposure , from 10nM to 10mM , and control cells were disposed in parallel with vehicle dimethyl sulfoxide.	PC9		
Significant circRNA , miRNA and mRNA alterations between PC9 and PC9ER cells were detected using limma packages and when fold change (FC)>=1.5 , as the selected FC cutoff .	PC9ER		
Software and Algorithms			
After purification , the labeled RNAs were hybridized to the microarrays , and the resulting arrays were scanned using an Agilent Scanner G2505C	Agilent Scanner		
Further , the raw data were normalized using the quantile algorithm , limma packages in R .	limma		Suggestion: LIMMA (RRID:SCR_010943)(link)
CeRNA Network Construction The circRNA-miRNA interactions were predicted using the Arraystars homemade miRNA target prediction software based on TargetScan (http:www.targetscan.org) and miRanda , and the R package miRNA tap was used to predict the target mRNAs .	TargetScan		
Five databases were included in the analysis , including PicTar , DIANA , TargetScan , miRanda and miRDB , wherein miRNA-mRNA interactions were reported consistently in at least two of the five databases .	PicTar		Suggestion: PicTar (RRID:SCR_003343)(link)
Cytoscape (v3.7.2) was used for network visualization , only including differentially expressed circRNAmiRNAmRNA with log2 FC >=1.5 in the analysis .	Cytoscape		Suggestion: Cytoscape (RRID:SCR_003032)(link)
2.5 PPI Network Construction and hub-Genes Identification The PPI network was built with the ceRNA network using the STRING program , version 11.0 .	STRING		Suggestion: STRING (RRID:SCR_005223)(link)
The enriched GO and KEGG terms (adjusted p-value<.	KEGG		

All figures were analyzed using GraphPad Prism 7 (GraphPad Software, La Jolla, CA, USA).	GraphPad Prism		
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Two-group differences were determined using t-test, and statistical significance was set at P <.	t-test
Oligonucleotides	
The PCR primers used in this study were as follows: hsa_circ_0007312, 5-TAACGGTGACTAATGGTGT TAAAGG-3 (forward) and 5-GATGCATTTACAAGCAAATCTG TGC-3(reverse) hsa_circ_0003748, 5-GAGAAGGCAGCCAACCAGAT-3 (forward) and 5-TCTTCTTCTAACTTATGGCTGC ATG-3 (reverse) hsa_circ_0001398, 5-TGAGCTGATCAAGTTGACAGCA TC-3 (forward) and 5-CAACTCAATAAAC TTCAAGTACCGA-3 (reverse) hsa-miR-764, 5-GCAGGTGCTCACTTGTCCTCC T-3 (forward) and 5-GCGAGCACAGAATTAATACGAC -3 (reverse) hsa-miR-628 to 5p, 5-ATGCTGACATATTTACTAGAGG -3 (forward) and 5-GCGAGCACA GAATTAATACGAC-3 (reverse) MAPK1, 5-TGTTCCCAAATGCTGACT-3 (forward) and 5-AACTTGAATGGTGCTTCG -3 (reverse).	5-CAACTCAATAAAC TTCAAGTACCGA-3

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq