Document Identifier: **PMC8659406**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Primary fibroblasts were isolated from surgically discarded foreskin samples obtained by an exempt Institutional Review Board protocol approved by the Duke University (Durham , NC) .
Consent: Patient consent for experiments was not required because the United States laws consider human tissue left over from surgery as discarded material.
IACUC: Animal studies Animal studies were performed in compliance with Duke Animal Care and Use Committee.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication

4,5-dimethylthiazol-2-yl)-2,5 diphenyl tetrazolium bromide assay , melanoma cells (A375 and A2058) , SCC cells (A431 , CAL27 , and Fadu) , and primary human dermal fibroblasts were seeded at 8 x 103 cells per well in a 96-well plate , and after 24 hours , cells were treated in triplicates with increasing concentrations of DLAD or LLAD for 24 , 48 , and 72 hours .
Number: 4,5-dimethylthiazol-2-yl)-2,5 diphenyl tetrazolium bromide assay , melanoma cells (A375 and A2058) , SCC cells (A431 , CAL27 , and Fadu) , and primary human dermal fibroblasts were seeded at 8 x 103 cells per well in a 96-well plate , and after 24 hours , cells were treated in triplicates with increasing concentrations of DLAD or LLAD for 24 , 48 , and 72 hours .
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Protein analysis by immunostaining For immunofluorescent staining, 5-6 mm thick cryosections were fixed in cold 100 methanol for about 20 minutes, blocked with 10 horse serum, incubated with primary antibodies against Ki-67 (MA5-14520, Thermo Fisher Scientific), CD45 (Sc-53665, Santa Cruz Biotechnology, Dallas, TX), and F480 (123103, BioLegend, San Diego, CA), followed by detection with Florescent Dylight-555-conjugated secondary antibodies and counterstained with DAPI (Thermo Fisher Scientific).	Ki-67		
	CD45		
	Sc-53665, Santa Cruz Biotechnology, Dallas, TX		
Experimental Models: Cell Lines			
Subcutaneous tumors were generated by injection of A2058 and A375 human melanoma cells (20,000 cells in 100 ml PBS-to-matrigel 3:1	A375		Suggestion: RRID:CVCL_0132(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The statistical difference in means between groups was evaluated in R Foundation for Statistical Computing (R Foundation, Vienna, Austria) using a two-tailed Students t-test unless specified otherwise, and P < 0.05 was considered statistically significant.	Students t-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq