Document Identifier: **PMC8664339**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: 15 All kidneys were donated by relatives or deceased citizens, and all kidneys were donated voluntarily with written informed consent, which was conducted in accordance with the Declaration of Istanbul.
IRB: This study was approved by our institutional Ethics Review Committee and was conducted in accordance with the principles of the Declaration of Helsinki.
Inclusion and Exclusion Criteria
Adult CRKP-infected recipients (≥18 years of age) receiving CAZ-AVI for ≥72 h and other antibiotic-treated CRKP-infected recipients were included in the study, while recipients with colonized or contaminated culture results were excluded.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: Adult CRKP-infected recipients (≥18 years of age) receiving CAZ-AVI for ≥72 h and other antibiotic-treated CRKP-infected recipients were included in the study, while recipients with colonized or contaminated culture results were excluded.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Minimum inhibitory concentrations were interpreted according to breakpoints established by the Clinical and Laboratory Standards Institute.19 CRKP was defined as insensitivity to at least one carbapenem, with a minimum inhibitory concentration ≥2 mg/mL for ertapenem and ≥4 mg/mL for imipenem or meropenem.19 Statistical Analysis Statistical analysis was performed using SPSS software (version 25.0; SPSS Inc., Chicago, IL).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The independent-sample t tests and Mann–Whitney U-test were used to compare normal and non-normal continuous variables, respectively.	t tests
	Mann–Whitney U-test
Categorical variables were represented by frequency and percentage. Chi-square or Fischer exact tests were used for comparison of categorical variables between groups, as appropriate.	Chi-square
	Fischer exact tests
Kaplan–Meier analysis and Log rank test were used to evaluate differences in the 30-day survival curves of CRKP-infected recipients treated with or without CAZ-AVI.	Log rank test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq