Document Identifier: **PMC8665688**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: This study was performed with the approval of the Ethics Committee of Kumamoto University (approval number: 174) and in accordance with the Good Clinical Practice and the Declaration of Helsinki guidelines .
Inclusion and Exclusion Criteria
We excluded human papillomavirus (HPV)-positive tumours from the analysis-based immunostaining results for p16 , a surrogate marker for HPV infection .
Attrition
2.17 miRNA screening with 3D-gene microarray From the advanced OSCC patients who received chemoradiotherapy , serum samples were obtained from three patients who responded to chemoradiotherapy and four who did not (Table S4) .
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Each sample was run in triplicate.

Number: Each sample was run in triplicate.
Cell Line Authentication
Authentication: 2.2 Short tandem repeat analysis Genomic DNA was extracted from SAS and SAS-R using NucleoSpin Tissue kit (MACHEREY-NAGEL, Duren, Germany), and short tandem repeat (STR) analysis of the extracted gDNA was performed by Takara Bio (Shiga, Japan).

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The antibodies used were mouse anti-CD9 antibody (12A12 COSMO BIO) ,	anti-CD9		
11,000 mouse anti-CD81 antibody (12C4 COSMO BIO) ,	anti-CD81		
11,000 mouse anti-ALIX antibody (ab117600 Abcam , Cambridge , UK)	anti-ALIX		
1500 mouse anti-GAPDH antibody (ab8245 Abcam) ,	anti-GAPDH		
11,000 rabbit anti-calnexin antibody (2679 Cell Signalling Technology , Danvers , MA , USA)	anti-calnexin		
11,000 anti-mouse IgG , HRP-linked antibody (7076 Cell Signalling Technology) , 12,000 and anti-rabbit IgG , HRP-linked antibody (7074 Cell Signalling Technology) , 12,000	anti-mouse IgG		
	HRP-linked antibody (7076 Cell Signalling Technology)		
	anti-rabbit IgG		
	HRP-linked antibody (7074 Cell Signalling Technology)		
Experimental Models: Cell Lines			
2 MATERIALS AND METHODS 2.1 Cell lines and culture The human OSCC cell lines, SAS, and HSC-2 were purchased from the Japanese Collection of Research Bioresources Bank of the National Institutes of Biomedical Innovation, Health and Nutrition (Osaka, Japan).	HSC-2		Suggestion: RRID:CVCL_1287(link)
Recombinant DNA			
After 24 h , pCT-CD9-GFP (pCMV , ExosomeSecretory	pCT-CD9-GFP		
	pCMV		
In the same way , pCT-CD9-RFP (SBI) was transfected into SAS-R cells .	pCT-CD9-RFP		
Software and Algorithms			
The videos were recorded and analyzsd with NTA software version 3.3 to determine the size and concentration of the particles . 2.8 Uptake of EVs SAS EVs or	NTA		

SAS-R EVs (2 mg) were stained with the red fluorescent dye PKH26 (Sigma-Aldrich) according to the manufacturers instructions .			
The relative expression of each protein was calculated using the Image J 1.52q software (National Institutes of Health , Bethesda , MD , USA) .	Image J		Suggestion: ImageJ, RRID:SCR_003070(link)
All analyses were performed with JMP-9 software (SAS Institute Inc, Cary, NC, USA).	SAS Institute		Suggestion: Statistical Analysis System, RRID:SCR_008567(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The Chi-squared test was performed to determine the associations between the miR-503-3p expression status and clinical or pathological variables .	Chi-squared test
The Kaplan-Meier method was used to estimate the probability of OS and DFS as a function of time , and the statistical differences in the survival of the subgroups of patients were compared by the log-rank test .	log-rank test
2.22 Statistical analysis Differences in mean values between two groups were analysed using Students t-tests, while differences in mean values among multiple groups were analysed by one-way ANOVA with the BonferroniDunn test.	t-tests
	ANOVA
	BonferroniDunn test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq