Document Identifier: **PMC8693313**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Materials and methods Ethics The use of human tissue samples and animal studies was approved by the Clinical Research Committee and the Institutional Animal Care and Use Committee of the General Hospital of Northern Theater Command, respectively.
IACUC: Materials and methods Ethics The use of human tissue samples and animal studies was approved by the Clinical Research Committee and the Institutional Animal Care and Use Committee of the General Hospital of Northern Theater Command, respectively.
Consent: All human studies adhered to the guidelines of the Declaration of Helsinki, and all participants provided informed consent.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
7 Animals Male C57BL/6 mice (8–10 weeks old) were purchased from the Nanjing Model Animal Center (Nanjing, China).
Subject Demographics
Age: Patients above 18 years of age, with or without T2DM and recommended and implanted with an implantable cardioverter defibrillator (ICD) without heart failure as per the guidelines, were considered eligible.
Randomization
The mice were fed HFD (Research Diets, New Brunswick, NJ) for 12 weeks and then divided into two groups at random (n = 8 per group).
Blinding
not detected.
Power Analysis

not detected.
Replication
not required.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Coimmunoprecipitation of RNA with Ago2 antibody 3T3-L1 cells were lysed using IP buffer (Thermo Fisher Scientific) and then immunoprecipitated with an anti-Ago2 antibody (Abnova, Taiwan, China) or IgG (Santa Cruz Biotechnology, Santa Cruz, CA, USA) binding to the agarose beads with protein A.	anti-Ago2		
	IgG		
Experimental Models: Cell Lines			
Cell culture and treatments 3T3-L1 and AD293 cells were obtained from the Chinese Academy of Sciences Cell Bank.	AD293		Suggestion: RRID:CVCL_9804(link)
Preadipocyte differentiation medium (ScienCell Research Laboratories, Carlsbad, CA, USA) was used to induce 3T3-L1 cell differentiation into mature adipocytes. miRNA mimics (100 nM), miRNA inhibitors (100 nM), and their controls (RiboBio, Guangzhou, China) were transfected into 3T3-L1 cells using X-tremeGENE HP DNA Transfection Reagent (Sigma, St. Louis, MO, USA), following the manufacturer's protocol.	3T3-L1		Suggestion: RRID:CVCL_0123(link)
Experimental Models: Organisms/Strains			
7 Animals Male C57BL/6 mice (8–10 weeks old) were purchased from the Nanjing Model Animal Center (Nanjing, China).	C57BL/6		Suggestion: RRID:IMSR_JAX:000664(link)
Recombinant DNA			
To construct luciferase reporter plasmids, we inserted the 3# UTR and mutant 3# UTR of the HDAC9 gene into the pmiR-RB-Report (RiboBio).	pmiR-RB-Report		
The CDS region of mmu-miR-574 was inserted into the pcDNA3.1-tGFP vector (Promega, Beijing, China).	pcDNA3.1-tGFP		
The mmu-miR-574 promoter vector and the mutated promoter vector were cloned into the pGL3 plasmids.	pGL3		Suggestion: RRID:Addgene_44264(link)
Software and Algorithms			

Preadipocyte differentiation medium (ScienCell Research Laboratories, Carlsbad, CA, USA) was used to induce 3T3-L1 cell differentiation into mature adipocytes. miRNA mimics (100 nM), miRNA inhibitors (100 nM), and their controls (RiboBio, Guangzhou, China) were transfected into 3T3-L1 cells using X-tremeGENE HP DNA Transfection Reagent (Sigma, St. Louis, MO, USA), following the manufacturer's protocol.	Sigma		
The binding position of HDAC9 and miR-574-5p was predicted by Database: miRWalk, and the seed sequence of miR-574-5p targeted HDAC9 in its CDS regions in both humans and mice, as shown in Figure 3D.	miRWalk		Suggestion: miRWalk (RRID:SCR_016509)(link)
All statistical analyses were performed using the SPSS version 24.0.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Between-group comparisons were performed using Student’s t test, whereas one-way analysis of variance was used for comparisons of more than two groups.	t test
	analysis of variance

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq