Document Identifier: **PMC8693952**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The slides were then incubated with Alexa-conjugated secondary antibodies (Thermo Fisher Scientific) for 1 h at RT , washed , and mounted with Fluoro-KEEPER Antifade Reagent (Nacalai Tesque)	Fluoro-KEEPER Antifade Reagent (Nacalai Tesque)		
For detection , the slides were washed with Tris-buffered saline containing 0.1 Tween-20 (TBST) , incubated with a blocking solution (3 BSA prepared in TBST) at RT for 1 h , and subsequently incubated with anti-DIG antibodies diluted in the blocking solution at 4degC overnight .	anti-DIG		
Experimental Models: Cell Lines			
Cell culture We generated HCT116TR_DBC1-3xFLAG cells using the Flp-In System (Thermo Fisher Scientific) .	HCT116TR_DBC1-3xFLAG		
First , we generated the doxycycline-induced HCT116FRT stable cell lines , HCT116FRTTR , that express the tetracycline repressor (TR) from pcDNA6TR-IRES-puro .	HCT116FRT		
HCT116TR_DBC1-3xFLAG cells were prepared using the isolated HCT116FRTTR by the Flp-In System and cultured at 37degC with 5 CO2 in DMEM10 fetal bovine serum (FBS) , supplemented with 2 mgml puromycin and 200 mgml hygromycin B . HCT116 , NIH3T3 , and HeLa cells were cultured in DMEM10 FBS at 37degC with 5 CO2 .	HCT116FRTTR		
	NIH3T3		Suggestion: RRID:CVCL_0594(link)
	HeLa		Suggestion: RRID:CVCL_0030(link)
The supernatant containing the HCT116 cells extract was mixed with the anti-DYKDDDDK magnetic agarose (Thermo Fisher Scientific) and rotated at 4degC overnight .	HCT116		Suggestion: RRID:CVCL_0291(link)
Recombinant DNA			
Plasmid construction The pcDNA5FRTTO-DBC1-3xFLAG plasmid was generated by inserting the PCR-amplified DBC1-3xFLAG sequence into the pcDNA5FRTTO	pcDNA5FRTTO-DBC1-3xFLAG		
	pcDNA5FRTTO		Suggestion: RRID:Addgene_135107(link)

vector (Thermo Fisher Scientific) between the EcoRV and XhoI sites .			
The pcDNA6TR-IRES-puro plasmid was constructed by ligating the PCR-amplified sequence containing a pUC origin to the TetR gene of pcDNA6TR (Thermo Fisher Scientific) with the segment of IRES to the puromycin-resistance gene of pCAGGS-FLPe (Gene Bridges) between the EcoRI and PstI sites .	pcDNA6TR		
	pCAGGS-FLPe		Suggestion: RRID:Addgene_20733(link)
The pMAL_TEV-HNRNPL plasmid was generated by inserting the PCR-amplified HNRNPL sequence into the pMAL_TEV vector (Yoshizawa et al. , 2018) between the HindIII and BamHI sites .	pMAL_TEV-HNRNPL		
	pMAL_TEV		
First , we generated the doxycycline-induced HCT116FRT stable cell lines , HCT116FRTTR , that express the tetracycline repressor (TR) from pcDNA6TR-IRES-puro .	pcDNA6TR-IRES-puro		
Software and Algorithms			
Protein identification was performed using ProteinPilot software 2.0 (AB Sciex) against the human UniProt database (version 2015_11) as described by Tohsato et al. (2012) .	ProteinPilot		
	UniProt		
The nucleoplasm and DNB volumes as well as the proportion of HNRNPL protein present within the DNB were estimated from the IF intensity profile over different Z positions (Z-stack) using cellSens Dimension software (Olympus) .	cellSens		
Protein concentrations were determined by absorbance at 280 nm using their extinction coefficients predicted by the ProtParam tool .	ProtParam		

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq