Materials Design Analysis Reporting (MDAR) Checklist for Authors

The MDAR framework establishes a minimum set of requirements in transparent reporting applicable to studies in the life sciences (see Statement of Task: doi:10.31222/osf.oio/9sm4x.). The MDAR checklist is a tool for authors, editors and others seeking to adopt the MDAR framework for transparent reporting in manuscripts and other outputs. Please refer to the MDAR Elaboration Document for additional context for the MDAR framework.

Materials

Antibodies	Yes (indicate where provided: page no/section/legend)	n/a
For commercial reagents, provide supplier name, catalogue number and RRID, if available	No antibodes detected. Please add identifiers for all resources where possible	

Cell Materials	Yes (indicate where provided: page no/section/legend)	n/a
Cell lines: Provide species information, strain. Provide accession number in repository OR supplier name, catalog number, clone number, OR RRID	No cell lines detected Please add identifiers for all resources where possible	
Primary cultures: Provide species, strain, sex of origin, genetic modification status.	Not currently checked by SciScore	

Experimental Animals	Yes (indicate where provided: page no/section/legend)	n/a
Laboratory animals: Provide species, strain, sex, age, genetic modification status. Provide accession number in repository OR supplier name, catalog number, clone number, OR RRID	No organisms detected Please add identifiers for all resources where possible	
Animal observed in or captured from the field: Provide species, sex and age where possible	Not currently checked by SciScore	
Model organisms: Provide Accession number in repository (where relevant) OR RRID	See laboratory animals section for information.	

Plants and microbes	Yes (indicate where provided: page no/section/legend)	n/a
Plants: provide species and strain, unique accession number if available, and source (including location for collected wild specimens)	Not currently checked by SciScore	
Microbes: provide species and strain, unique accession number if available, and source	Not currently checked by SciScore	

Human research participants	Yes (indicate where provided: page no/section/legend)	n/a
Identify authority granting ethics approval (IRB or equivalent committee(s), provide reference number for approval.	The local ethics committee of the University of Erlangen-Nuremberg approved the study protocol.	
Provide statement confirming informed consent obtained from study participants.	Each patient provided written informed consent prior to study inclusion.	
Report on age and sex for all study participants.	**Age:**Study population Individuals, aged between 18 and 85 years, were included in the study, if they had a HF with reduced (HFrEF) or mid-range ejection fraction (HFmrEF), according to the ESC guidelines definition, in stable condition New York Heart Association (NYHA) II-III. 12 Patients could have diabetes or not however, key exclusion criteria were any other form of diabetes than type 2 diabetes, the use of insulin, or any SGLT2 inhibitor 10 weeks prior to inclusion. **Sex:**not detected.	

Design

Study protocol	Yes (indicate where provided: page no/section/legend)	n/a
For clinical trials, provide the trial registration number OR cite DOI in manuscript.	NCT03128528	

Laboratory protocol	Yes (indicate where provided: page no/section/legend)	n/a
Provide DOI or other citation details if detailed step-by-step protocols are available.	Not detected.	

Experimental study design (statistics details)	Yes (indicate where provided: page no/section/legend)	n/a
State whether and how the following have been done, or if they were not carried out		
Sample size determination	not detected.	
Randomization	Methods Study design This was an investigator initiated, double-blind, randomized, placebo-controlled, parallel-group, prospective phase II single centre study, performed at the Clinical Research Unit (CRC) of the Department of Nephrology and Hypertension, University Hospital Erlangen-Nuremberg, Germany, in order to analyse changes of vascular function and of tissue sodium content in patients with HF after SGLT2 inhibition with empagliflozin.	
Blinding	not detected.	
inclusion/exclusion criteria	Study population Individuals, aged between 18 and 85 years, were included in the study, if they had a HF with reduced (HFrEF) or mid-range ejection fraction (HFmrEF), according to the ESC guidelines definition, in stable condition New York Heart Association (NYHA) II-III. 12 Patients could have diabetes or not however, key exclusion criteria were any other form of diabetes than type 2 diabetes, the use of insulin, or any SGLT2 inhibitor 10 weeks prior to inclusion.	

Sample definition and in-laboratory replication	Yes (indicate where provided: page no/section/legend)	n/a
State number of times the experiment was replicated in laboratory	Not detected.	
Define whether data describe technical or biological replicates	Not detected.	

Ethics	Yes (indicate where provided: page no/section/legend)	n/a
Studies involving human participants: State details of authority granting ethics approval (IRB or equivalent committee(s), provide reference number for approval.	The local ethics committee of the University of Erlangen-Nuremberg approved the study protocol.	
Studies involving experimental animals: State details of authority granting ethics approval (IRB or equivalent committee(s), provide reference number for approval.	Not detected.	
Studies involving specimen and field samples: State if relevant permits obtained, provide details of authority approving study; if none were required, explain why.	Not detected.	

Dual Use Research of Concern (DURC)	Yes (indicate where provided: page no/section/legend)	n/a
If study is subject to dual use research of concern, state the authority granting approval and reference number for the regulatory approval	Not currently checked by SciScore	

Analysis

Attrition	Yes (indicate where provided: page no/section/legend)	n/a
State if sample or data point from the analysis is excluded, and whether the criteria for exclusion were determined and specified in advance.	not detected.	

Statistics	Yes (indicate where provided: page no/section/legend)	n/a
Describe statistical tests used and justify choice of tests.	Statistical significance of changes between baseline and 1 or 3 months treatment with empagliflozin or placebo were determined using paired t-test, whereas for differences between 1 or 3 months treatment with empagliflozin compared with placebo, an unpaired t-test was used.	

Data availability	Yes (indicate where provided: page no/section/legend)	n/a
State whether newly created datasets are available, including protocols for access or restriction on access.	Not detected.	
If data are publicly available, provide accession number in repository or DOI or URL.	Not detected.	
If publicly available data are reused, provide accession number in repository or DOI or URL, where possible.	Not detected.	

Code availability	Yes (indicate where provided: page no/section/legend)	n/a
For all newly generated code and software essential for replicating the main findings of the study:		
State whether the code or software is available.	Not detected.	
If code is publicly available, provide accession number in repository, or DOI or URL.	Not detected.	

Analysis

Adherence to community standards	Yes (indicate where provided: page no/section/legend)	n/a
MDAR framework recommends adoption of discipline-specific guidelines, established and endorsed through community initiatives. Journals have their own policy about requiring specific guidelines and recommendations to complement MDAR.		
State if relevant guidelines (eg., ICMJE, MIBBI, ARRIVE) have been followed, and whether a checklist (eg., CONSORT, PRISMA, ARRIVE) is provided with the manuscript.	Not currently checked by SciScore	