Document Identifier: **PMC8724682**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: All donors provided written informed consent, and the investigators were not able to ascertain the identity of the donors.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
2.2 2D model 2.2.1 hGF cell culture Primary hGFSS (27 years, Caucasian, female, lot number 313 x 100401) were used.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
2.2.2 Treatment with the MIM A blind study was performed using the vehicle and two treatments containing LD BMP2 and BMP4.
Power Analysis
not detected.
Replication
The experiment was run in six sample replicates (n = 6) for each group.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The sections were incubated overnight at 4degC with mouse monoclonal primary antibodies (Table 1) treated with biotinylated anti-mouse secondary antibody+++++++ +++++++ for 30 minutes at RT and followed by incubation in avidin-biotinylated peroxidase complex for 30 minutes at RT.	anti-mouse		
TABLE 1 Primary antibodies used for immunohistochemical staining Antibody Dilution Clone Isotype Keratin 17 a 1:50 E-4 IgG1 Keratin 19 a 1:50 A-3 IgG1 Involucrin a 1:20 SY5 IgG1 Vimentin a 1:1000 E-5 IgG1 a Santa Cruz Biotechnology Inc., Santa Cruz, CA.	E-4 IgG1		
	A-3 IgG1		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The Shapiro-Wilk test was done to assume parametric or non-parametric distribution for the normality tests.	Shapiro-Wilk test
Differences between groups were assessed, depending on their normal distribution, by the Mann-Whitney U test or the two-way analysis of variance (ANOVA) test, followed by post-hoc pairwise comparisons using the LSD test.	Mann-Whitney U test
	analysis of variance
	ANOVA
	LSD test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq