Document Identifier: **PMC8728431**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: https://www.flowjo.com/ Other Patient cell study Study was approved by the University of North Carolina Institutional Review Board UNC IRB committee Resource availability Lead contact Further information and requests for resources and reagents should be directed to the Lead Contact , Dr. Guochun Jiang (Guochun_Jiang@med.unc . edu) .
Consent: All participants provided informed consent and the study was approved by the UNC Institutional Review Board.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication

For the HIV gag RNA measurements in resting CD4+ T cells from ART-suppressed individuals , 3–5 replicates of 2–4 million resting CD4+ T cells per replicate were treated with compounds either for 24 or 48 hours .
Type: Three biological replicates were prepared for this experiment .
Number: For the HIV gag RNA measurements in resting CD4+ T cells from ART-suppressed individuals , 3–5 replicates of 2–4 million resting CD4+ T cells per replicate were treated with compounds either for 24 or 48 hours .
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
STAR#Methods Key resources table REAGENT or RESOURCE SOURCE IDENTIFIER Antibodies anti-cIAP1 abcam Cat#ab108361 anti-cIAP2 abcam Cat#ab23423 anti-USP7 Bethyl Cat#A700-072	anti-cIAP1		
	anti-cIAP2		
Clarified lysates were precleared with protein A/G agarose beads (Thermo Scientific) , followed by IP with either anti-USP7 monoclonal antibody (Bethyl) , anti-TRIM27 polyclonal antibody (MyBioSource) , or normal rabbit IgG (Cell Signaling) .	anti-USP7		
	anti-TRIM27		
	normal rabbit IgG		
Whole cell protein extracts were prepared with RIPA buffer (Sigma) followed by Co-IP with IgG or anti-PanCrotonylLysine antibody (PTM biolabs) .	anti-PanCrotonylLysine		
Experimental Models: Cell Lines			
models: Cell lines Jurkat ATCC TIB-152 2D10 Dr.	Jurkat		
J-Lat A1 cells , J-Lat 10.6 , or U1 cells were from NIH AIDS Reagents Repository , 2D10 cell model of latency was a gift from Dr. Jonathan Karn (Case Western University , Cleveland , OH) , which harbors a single copy of HIV-1pNL4-3 whose genome lacks sequences encoding the majority of the Gag-Pol polyprotein , but encompasses the full-length 5 and 3# LTRs .	J-Lat A1		
Software and Algorithms			
Trim 27 Accell small interfering RNAs Dharmacon Cat#E-006552-00-0020 scrambled control siRNAs Dharmacon Cat#D-001910-10-20 Software and algorithms GraphPad Prism 8 GraphPad Software Inc .	GraphPad Prism		
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)
https://www.graphpad.com/	https:// www.graphpad.com/		
The glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and TBP primers/probe sets were used for control PCR (Applied Biosystems Inc .) HIV (GFP) was quantified by	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)

GFP expression using flow cytometry and the data were analyzed using FlowJo Software.			
Whole cell protein extracts were prepared with RIPA buffer (Sigma) followed by Co-IP with IgG or anti-PanCrotonylLysine antibody (PTM biolabs) .	Sigma		
MS data analysis Mass spectra processing and peptide identification were performed on the Andromeda search engine in MaxQuant software (Version 1.6.0.16) against a human UniProt database (UP000005640) .	MaxQuant		Suggestion: MaxQuant (RRID:SCR_014485)(link)
	UniProt		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Protein quantitation was performed on biological replicate runs, and a two-sample T-test statistics was used to report statistically significant expression fold-changes.	T-test
Statistical significance was determined using the two-tailed Student T test or One-way ANOVA test, where ∗, p <0.05 was considered significant.	T test
	ANOVA
Oligonucleotides	
2018 Oligonucleotides HIV long LTR probe: 6FAMCCA GAG TCA CAC AAC AGA CGG GCA CAT AMRA Invitrogen Yukl ,	6FAMCCA GAG TCA CAC AAC AGA CGG GCA CAT AMRA
et al. 2018 HIV long LTR sense: GCC TCA ATA AAG CTT GCC TTG A Invitrogen Yukl ,	GCC TCA ATA AAG CTT GCC TTG
et al. 2018 HIV long LTR antisense: GGG CGC CAC TGC TAG AGA Invitrogen Yukl , et al. 2018 GAPDH (Hs99999905_m1) Thermo Scientific Cat#4331182 TRIM27 Applied Biosystems Hs01100632_ml	GGG CGC CAC TGC TAG AGA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq