Document Identifier: **PMC8743587**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The Institutional Review Board of the Seoul National University Hospital approved the storage of blood samples for genetic analysis with informed consent (IRB No. H-1103-127-357), and the Institutional Review Board approved this study protocol (IRB No. 1601-063-734) for the GENIE cohort.
Consent: The Institutional Review Board of the Seoul National University Hospital approved the storage of blood samples for genetic analysis with informed consent (IRB No. H-1103-127-357), and the Institutional Review Board approved this study protocol (IRB No. 1601-063-734) for the GENIE cohort.
Inclusion and Exclusion Criteria
Among 12,642 participants from the three cohorts, 4,856 were excluded, and 7,786 were enrolled.
Attrition
Among 12,642 participants from the three cohorts, 4,856 were excluded, and 7,786 were enrolled.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication

Replication analysis, rare variant analysis, and expression analysis For replication analysis, we searched the National Human Genome Research Institute-European Bioinformatics Institute GWAS catalog (https:www.ebi.ac.ukgwashome, December 2020 EFO_0004631) for SNPs, and selected 406 SNPs.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The GENIE cohort consists of mostly healthy people who undergo regular check-ups in the healthcare system.	GENIE		Suggestion: GENIE (RRID:SCR_009197)(link)
Biochemical measurements Serum 25(OH)D levels were measured by radioimmunoassay (DiaSorin Inc., Stillwater, MN, USA) for the GENIE cohort and by chemiluminescent microparticle immunoassay (CMIA) using an Architect i2000SR system (Abbott, Singapore) for the KARE and VHSMC cohorts.	Abbott		
The genome-wide significance level for SNPs was set at 5x10-8, annotated with ANNOVAR (ANNOtate VARiation) 34 and used for regional plots.	ANNOVAR		Suggestion: ANNOVAR (RRID:SCR_012821)(link)
We used PLINK 1.9 to evaluate the linkage disequilibrium (LD) across the ancestral population with Phase 3 haplotype data of 1000 Genome Projects.	PLINK		Suggestion: PLINK (RRID:SCR_001757)(link)
Moreover, we annotated genes by constructing gene interaction networks with the STRING v.	STRING		

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq