Document Identifier: **PMC8747765**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
The Kaplan–Meier (K-M) survival curve described the predictive power of indicators using the “survival” and “survminer” packages .
Replication
not required.
Cell Line Authentication
not detected.

Data Information	
Identifiers: GCH1 Marker 123 PTGS2 , DUSP1 , NOS2 , NCF2 , MT3 , UBC , ALB , TXNRD1 , SRXN1 , GPX2 , BNIP3 , OXSR1 , SELENOS , ANGPTL7 , CHAC1 , SLC7A11 , DDIT4 , LOC284561 , ASNS , TSC22D3 , DDIT3 , JDP2 , SESN2 , SLC1A4 , PCK2 , TXNIP , VLDLR , GPT2 , PSAT1 , LURAP1L , SLC7A5 , HERPUD1 , XBP1 , ATF3 , SLC3A2 , CBS , ATF4 , ZNF419 , KLHL24 , TRIB3 , ZFP69B , ATP6V1G2 ,	ATP6V1G2
Identifiers: Immunity Analysis The CIBERSORT,20,21 ESTIMATE,22 MCPcounter,23 single-sample gene set enrichment analysis (ssGSEA),24 and TIMER25 algorithms were used to assess cellular components or cell immune responses.	CIBERSORT,20,21

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
, IREB2 , ATP5MC3 , CS , EMC2 , ACSF2 , NOX1 , CYBB , NOX3 , NOX4 , NOX5 , DUOX1 , DUOX2 , G6PD , PGD , VDAC2 , PIK3CA , FLT3 , SCP2 , TP53 , ACSL4 , LPCAT3 , NRAS , KRAS , HRAS , TF , TFRC , TFR2 , SLC38A1 , SLC1A5	EMC2		
ELAVL1 , BAP1 , TP53 , ABCC1 , ACSL4 , MIR6852 , ACVR1B , TGFBR1 , BAP1 , EPAS1 , HILPDA , HIF1A , ALOX12 , ACSL4 , HMOX1 , IFNG , ANO6 , LPIN1 , HMGB1 , TNFAIP3 , TLR4 , NOX4 , ATF3 , ATM , YY1AP1 , EGLN2 , MIOX , TAZ , MTDH , IDH1 ,	ELAVL1		
EIF2S1 , KIM-1 , IL6 , CXCL2 , RELA , HSD17B11 , AGPAT3 , SETD1B , HMOX1 , TF , FTL , RPL8 , ATP5MC3 ,	EIF2S1		
Chemotherapy Sensitivity Analysis We obtained the drug sensitivity data and RNA-seq data of the NCI-60 cell line from CellMiner (https://discover.nci.nih.gov/ cellminer/home.do) and analyzed the relationship between the prognostic lncRNAs and drug sensitivity using Pearson correlation analysis.	NCI-60		
Software and Algorithms			
The Wilcoxon signed-rank test was utilized to identify differentially expressed ferroptosis-related genes and differentially expressed ferroptosis-related lncRNAs by the limma package of R based on the cutoff values: |Log2FC| > 1 and FDR < 0.05 .	limma		
The “clusterProfiler” package was used to perform Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses on differentially expressed ferroptosis-related genes.	KEGG		
Protein–Protein Interaction Network Construction The String database (https://cn.string-db.org/) was used to analyze the interactions	String		

of ferroptosis-related genes and ferroptosis-related lncRNAs .			
Statistical Analysis R version 4.0.2 software and its resource packages were used for statistical analysis and related visualization graphics.	Statistical Analysis R		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
HNF4A , NGB , YWHAE , GABPB1 , AURKA , MIR4715 , RIPK1 , PRDX1 , MIR30B Identification and Differential Expression Analysis of Ferroptosis-Related lncRNAs Pearson correlation analysis between the ferroptosis-related genes and lncRNAs was used to identify the ferroptosis-related lncRNAs (| R| > 0.5 and p < 0.001) .	Pearson correlation
The Wilcoxon signed-rank test was utilized to identify differentially expressed ferroptosis-related genes and differentially expressed ferroptosis-related lncRNAs by the limma package of R based on the cutoff values: |Log2FC| > 1 and FDR < 0.05 .	Wilcoxon signed-rank test
17,18 Development of the Ferroptosis-Related lncRNA Prognostic Signature Lasso penalized Cox regression and univariate Cox regression analysis were employed to construct the ferroptosis-related lncRNA prognostic signature , which was used to construct a risk scoring function .	Cox regression
Student's t test was used to compare differences between different risk groups.	t test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq