Document Identifier: **PMC8748218**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: This study was approved by the Ethics Committee of Hyogo College of Medicine (Rinhi-273), and performed in accordance with the Declaration of Helsinki (1995) of the World Medical Association (as revised in 2013 in Fortaleza, Brazil).
Consent: All patients provided written informed consent.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The aligned sequence files were sorted and merged using SAMtools (samtools.sourceforge.net, version 0.1.18).	samtools		
GATK (http:www.broadinstitute.orggatk) was used for realignment, base quality score recalibration, single nucleotide variant or indel (small insertions and deletions) variant calling, and variant quality recalibration.	GATK		Suggestion: GATK (RRID:SCR_001876)(link)
Germline or somatic mutations were confirmed by comparison between genomic DNA from tumors and that from the PBL of the patient, using Sanger sequencing.	Germline		Suggestion: GERMLINE (RRID:SCR_001720)(link)
JMP Pro v14.2 (SAS Institute, Cary, NC, USA) was used for the statistical analysis.	SAS Institute		Suggestion: Statistical Analysis System, RRID:SCR_008567(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
13 2.5 Statistical analyses Gene expression ratios were compared between, in ccRCCs, with 14q loss and the ones without 14q loss using the chi-squared test.	chi-squared test
Nonparametric pairwise multiple comparison analysis among 3 metastasis groups was carried out using Dunn multiple comparison test with the number of chromosome alterations.	Dunn multiple comparison test
Oligonucleotides	
The following primers were also used , after which normalization was conducted with glyceraldehyde-3-phosphate dehydrogenase (GAPDH) HIF1A: 5-CACCACAGGACAGTACAGGAT- 3 and 5-CGTGCTGAATAATACCACTCAC A-3 , L2HGDH: 5-TGTGCAGGACTTTACTCAGACC -3 and 5-GTAATCTCCCCGGAATGGTACA -3 , KLHL33: 5-GCGTGGGACTGGTACGAAC-3 and 5-GGGTGTTGGAGTGACTGTAGA-3 , GAPDH: 5-GCACCGTCAAGGCTGAGAAC-3 and 5-TGGTGAAGACGCCAGTGGA-3 .	5-TGTGCAGGACTTTACTCAGACC -3
	5-GCGTGGGACTGGTACGAAC-3

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq