Document Identifier: **PMC8751882**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: All animal experimental procedures were approved by the Institutional Animal Care and Use Committee at Loma Linda University (8190039) and were compliant with the Animal Research: Reporting In Vivo Experiments (ARRIVE) guidelines.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Reagents The AHR antagonist, 6,2,4-trimethoxyflavone (TMF), AHR agonist, 3-methylcholanthrene (3-MC), dimethyl sulfoxide (DMSO), dextrose, and mannitol were from Sigma-Aldrich, MO. LSKL, Thrombospondin (TSP-1) inhibitor, was from MedChem Express, NJ. Animal Models and Experimental Design A total of 387 adult male CD-1 mice (weight 30-35 g, Charles River, Wilmington, MA) aged 8-10 weeks were used for the study.
Subject Demographics
Weight: Reagents The AHR antagonist, 6,2,4-trimethoxyflavone (TMF), AHR agonist, 3-methylcholanthrene (3-MC), dimethyl sulfoxide (DMSO), dextrose, and mannitol were from Sigma-Aldrich, MO. LSKL, Thrombospondin (TSP-1) inhibitor, was from MedChem Express, NJ. Animal Models and Experimental Design A total of 387 adult male CD-1 mice (weight 30-35 g, Charles River, Wilmington, MA) aged 8-10 weeks were used for the study.
Age: Reagents The AHR antagonist, 6,2,4-trimethoxyflavone (TMF), AHR agonist, 3-methylcholanthrene (3-MC), dimethyl sulfoxide (DMSO), dextrose, and mannitol were from Sigma-Aldrich, MO. LSKL, Thrombospondin (TSP-1) inhibitor, was from MedChem Express, NJ. Animal Models and Experimental Design A total of 387 adult male CD-1 mice (weight 30-35 g, Charles River, Wilmington, MA) aged 8-10 weeks were used for the study.
Randomization
Mice were randomly assigned to different experimental groups by generating random numbers.
Blinding

Information on experimental groups was blinded to researchers who performed surgeries, neurobehavioral assessments, Western blot, immunofluorescence staining, and data analysis.
Power Analysis
not detected.
Replication
not required.
Cell Line Authentication
Authentication: To Determine the Role of AHR on HE and BBB Function After Hyperglycemic ICH Sixty-six mice were assigned to the following 4 groups: Sham, ICH+dextrose, ICH+dextrose+AHR clustered regularly interspaced short palindromic repeat (CRISPR) knockout, and ICH+dextrose+knockout CRISPR Control.
Data Information
Availability: Methods Data Availability All data are available within the article and additional data can be acquired from the corresponding author.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains			
Reagents The AHR antagonist, 6,2,4-trimethoxyflavone (TMF), AHR agonist, 3-methylcholanthrene (3-MC), dimethyl sulfoxide (DMSO), dextrose, and mannitol were from Sigma-Aldrich, MO. LSKL, Thrombospondin (TSP-1) inhibitor, was from MedChem Express, NJ. Animal Models and Experimental Design A total of 387 adult male CD-1 mice (weight 30-35 g, Charles River, Wilmington, MA) aged 8-10 weeks were used for the study.	CD-1		Suggestion: RRID:IMSR_CRL:022(link)
Software and Algorithms			
Western blot bands were quantified as relative density of bands using Image J software (NIH, Bethesda, USA).	Image J		Suggestion: ImageJ, RRID:SCR_003070(link)
Statistical Analysis All data analysis was performed using GraphPad Prism 8.2.1 (GraphPad Software, CA) and data were expressed as mean+-SD.	GraphPad Prism		Suggestion: GraphPad Prism (RRID:SCR_002798)(link)
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Neurobehavioral Tests The modified Garcia neurological score, corner turn test, forelimb placement test, and Rotorod test were used to assess neurobehavioral function.	Rotorod test
Multiple comparisons were statistically analyzed with Kruskal-Wallis test followed by post hoc Dunns method or 2-way ANOVA followed by the post hoc Tukey method.	Kruskal-Wallis test
	ANOVA
	Tukey method

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq