Document Identifier: **PMC8753994**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Patients and Methods Ethical Considerations, Eligibility, and Study Design This prospective randomized, double-blind clinical controlled study was approved by the Research Ethics Committee, South Egypt Cancer Institute, Assiut University, Assiut, Egypt (ID: SECI-IRBIORG0006563no. 402, date 23102017, Head of the Committee Prof.
Consent: Written informed consent was obtained from all study participants.
Inclusion and Exclusion Criteria
Exclusion criteria included significant cardiac , respiratory , renal , central nervous system , or hepatic diseases pregnancy BMI >=30 kgm2 bleeding diathesis , allergy to study drugs , history of drug addiction , and patients with psychiatric illnesses that would interfere with the perception and assessment of pain .
Attrition
Twenty-four patients were enrolled in two groups of 12 patients each to receive a bilateral single-injection TAP block with 20 mL of bupivacaine 0.5 (Markyrene(r) Sigma-Tec , Egypt) (TAP group) or 20 mL of bupivacaine 0.5 combined with 1 ugkg dexmedetomidine (Precedex(r
Sex as a biological variable
not detected.
Subject Demographics
Age: The patients enrolled in this study were aged 18-60 years, with body mass index (BMI)
Randomization
Patients and Methods Ethical Considerations, Eligibility, and Study Design This prospective randomized, double-blind clinical controlled study was approved by the Research Ethics Committee, South Egypt Cancer Institute, Assiut University, Assiut, Egypt (ID: SECI-IRBIORG0006563no. 402, date 23102017, Head of the Committee Prof.
Blinding
The surgeon , attending anesthesiologists , data collection personnel , and patients were blinded to the group assignment .
Power Analysis

Using the G-Power calculator 3.1.9.718 for the determination of sample size, 12 patients were needed in the TAP-DEX group to study dexmedetomidine pharmacokinetics based on a priori analysis with a type I error of 0.05, power of 0.8, and effect size of 0.8.	
Replication	
not required.	
Cell Line Authentication	
Authentication: Continuous data were checked for normality by visual inspection of histograms and by the Shapiro-Wilk test.	
Protocol Information	
Identifiers: (Identifier: NCT03328299, Principal investigator: Assistant Prof.	NCT03328299

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Using the G-Power calculator 3.1.9.718 for the determination of sample size, 12 patients were needed in the TAP-DEX group to study dexmedetomidine pharmacokinetics based on a priori analysis with a type I error of 0.05, power of 0.8, and effect size of 0.8.	G-Power		Suggestion: G*Power, RRID:SCR_013726(link)
Statistical Tests Data were collected and processed using SPSS version 20 (SPSS Inc., Chicago, IL, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Continuous data were checked for normality by visual inspection of histograms and by the Shapiro-Wilk test.	Shapiro-Wilk test
Categorical data were expressed as numbers and frequencies () and analyzed using the chi-square or Fishers exact test, as appropriate.	chi-square
	Fishers exact test
For normally distributed data, Students t-test and paired Students t-test were used for the analysis of independent and paired samples, respectively.	t-test
For abnormally distributed and ordinal data, the Mann-Whitney U-test and Wilcoxon signed-rank test were used for the analysis of independent and paired samples, respectively.	Mann-Whitney U-test
	Wilcoxon signed-rank test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq