Document Identifier: **PMC8754515**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The RS was approved by the Medical Ethics Committee of Erasmus MC and by the Ministry of Health, Welfare, and Sport of the Netherlands.
Consent: All participants provided written informed consent to participate in the study and to have their information obtained from treating physicians.
Inclusion and Exclusion Criteria
Only the SNPs that were also GWAS significant in the UK Biobank (i.e. , P < 1.0 x 10-8) were considered for inclusion in the phosphate GRS.
Attrition
Serum leptin concentrations were determined during the third visit of RS-I , in a random subset of participants selected as part of a separate case-cohort study (n = 489 men and n = 694 women) .
Sex as a biological variable
RS is a population-based study of men and women aged 40 y or older and recruited in the district of Ommoord, Rotterdam (20).
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
MR, Mendelian randomization analyses RS, Rotterdam Study
Blinding
not detected.
Power Analysis
not detected.
Replication

not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
All analyses were performed with IBM SPSS software , version 21 (SPSS)	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)
To account for potential overestimation of results due to family relatedness, 2sls MR analyses were repeated after randomly excluding first- and second-degree relatives, estimated using KING software.	KING		Suggestion: KING (RRID:SCR_009251)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis Differences between men and women were compared using independent t test for continuous variables and kh2 test for categorical variables .	t test
Because fat mass and lean mass are related to one another , we tested the correlation between fat mass and lean mass using Spearman correlation coefficients .	Spearman correlation
To correct for multiple testing of the independence assumption, a Bonferroni correction was applied resulting in a corrected P value < 0.006 (0.059) testing age, sex, education level, smoking, total calcium, 25(OH)D, eGFR, testosterone, and 17b-estradiol.	Bonferroni

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq