Document Identifier: **PMC8761437**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
2 MATERIALS AND METHODS 2.1 Patient information Ten tumour specimens from 10 male patients with SLSCC diagnosed by biopsy and pathology were collected immediately after surgical tumour resection.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Number: 2.12 Statistical analysis Data were presented as the mean +- standard deviation (SD) of at least three independent assays and analysed by Students t test or Fishers exact test.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The coverslips were immersed in solutions containing primary antibodies (rabbit antihuman) for pan-cytokeratin (CK 1:400 Abcam), vimentin (1:200 Abcam), a-smooth muscle actin (a-SMA 1:200 Abcam) and fibroblast activation protein (FAP 1:250 Abcam) and incubated at 4degC overnight.	antihuman) for pan-cytokeratin		
	vimentin		
	a-smooth muscle actin		
	a-SMA		
	fibroblast activation protein		
	FAP		
The cells were then treated with secondary antibodies (Jackson ImmunoResearch), either fluorescein isothiocyanate (FITC)-conjugated goat anti-rabbit IgG (H + L 1:100) or Cy(tm)3-conjugated goat anti-rabbit IgG (1:100) for 1 h at 37degC.	anti-rabbit IgG		
The membranes were initially blocked with 5 nonfat milk and probed overnight with rabbit primary anti-CD63 (1:1000, Abcam), anti-TSG101 (1:2000, Abcam) and anti-Calnexin (1:2000, Abcam), followed by detection using horseradish peroxidase-conjugated goat anti-mouserabbitrat IgG (1:2000 Jackson).	anti-Calnexin		
Beta-actin antibody (1:5,000 HuaAn Biotech) was used to normalise the amount of sample loaded.	Beta-actin		
Software and Algorithms			
The genes targeted by selective exosomal miRNAs were predicted by four online bioinformatic algorithms: TargetScan (http:www.targetscan.orgvert_72), miRDB (http:mirdb.org), miRTarBase (http:mirtarbase.mbc.nctu.edu.twphpindex.php) and miRWalk (http:mirwalk.umm.uni-heidelberg.de).	TargetScan		
	miRWalk		Suggestion: miRWalk (RRID:SCR_016509)(link)
2.11 Gene Ontology (GO) and Kyoto Encyclopaedia of Genes and Genomes (KEGG) analysis The GO analysis (http:www.geneontology.org) was used to predict the potential	KEGG		
	http:www.geneontology.org		
	http:www.genome.jp		
	DAVID		Suggestion: DAVID (RRID:SCR_001881)(link)

functions of the target genes of selected miRNAs, while the KEGG (http:www.genome.jp) pathway enrichment analysis was performed to identify the predominant pathways based on the DAVID 6.7 online software (http:david.abcc.ncifcrf.govhome.jsp).			
All analyses were performed with the GraphPad Prism software version 6.00 for Windows (GraphPad Software).	GraphPad Prism		
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
All GO terms and KEGG pathway enrichment analysis were analysed using kobas3.0 software (hypergeometric testFishers exact test) with a $p < 0.05$ threshold of significance, and the FDR correction method was Benjamini and Hochberg (Appendix S4).	Benjamini and Hochberg
2.12 Statistical analysis Data were presented as the mean +- standard deviation (SD) of at least three independent assays and analysed by Students t test or Fishers exact test.	t test
	Fishers exact test
Oligonucleotides	
TABLE 1 Sequences of primers used for quantitative real-time PCR miRNA name Primer sequence (5-3) miR-16-5p-R CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGCGCCAATA miR-16-5p-S ACACTCCAGCTGGGTAGCAGCACG TAAATA miR-29a-3p-R CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGTAACCGAT miR-29a-3p-S ACACTCCAGCTGGGTAGCACCATC TGAAAT miR-34c-5p-R CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGGCAATCAG miR-34c-5p-S ACACTCCAGCTGGGAGGCAGTGTA GTTAGCT miR-32-5p-R CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGTGCAACTT miR-32-5p-S ACACTCCAGCTGGGTATTGCACAT TACTAA miR-490-5p-R CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGACCCACCT miR-490-5p-S ACACTCCAGCTGGGCCATGGATCT CCAG miR-193b-3p-R CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGAGCGGGAC miR-193b-3p-S ACACTCCAGCTGGGAACTGGCCCT CAAAGT miR-10a-3p-R CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGTATTCCCC miR-10a-3p-S ACACTCCAGCTGGGCAAATTCGTA TCTAGG miR-656-3p-R CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGAGAGGTTG miR-656-3p-S	CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGCGCCAATA
	ACACTCCAGCTGGGTAGCAGCACG TAAATA
	CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGTAACCGAT
	ACACTCCAGCTGGGTAGCACCATC TGAAAT
	CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGGCAATCAG
	ACACTCCAGCTGGGAGGCAGTGTA GTTAGCT
	CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGTGCAACTT
	ACACTCCAGCTGGGTATTGCACAT TACTAA
	CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGACCCACCT
	ACACTCCAGCTGGGCCATGGATCT CCAG
	CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGAGCGGGAC
	ACACTCCAGCTGGGAACTGGCCCT CAAAGT
	CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGTATTCCCC
	ACACTCCAGCTGGGCAAATTCGTA TCTAGG
	CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGAGAGGTTG
	ACACTCCAGCTGGGAATATTATAC AGTCA
	CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGTCAGTTTT
	ACACTCCAGCTGGGTGTGCAAATC CATGCAA
	CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGTTCAAAAC

ACACTCCAGCTGGGAATATTATAC AGTCA miR-19b-3p-R CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGTCAGTTTT miR-19b-3p-S ACACTCCAGCTGGGTGTGCAAATC CATGCAA miR-424-5p-R CTCAACTGGTGTCGTGGAGTCGGC AATTCAGTTGAGTTCAAAAC miR-424-5p-S ACACTCCAGCTGGGCAGCAGCAAT TCATGT URP TGGTGTCGTGGAGTCG U6-S CTCGCTTCGGCAGCACA U6-A AACGCTTCACGAATTTGCGT John Wiley & Sons, Ltd 2.10 Identification of candidate exosomal miRNAs and target genes The miRNAs that showed significant differences ($p < 0.05$) between CAFs and NFs were selected as candidate exosomal miRNAs for further analysis.	ACACTCCAGCTGGGCAGCAGCAAT TCATGT URP TGGTGTCGTGGAGTCG U6-S CTCGCTTCGGCAGCACA AACGCTTCACGAATTTGCGT

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq