Document Identifier: **PMC8762459**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: All other reagents, unless otherwise stated, were obtained from Sigma Aldrich Ltd. 2.2 Animals Animal care, procedures, and ethical approval were obtained from UCD, Ireland and UCC Ethics Committee.
Euthanasia Agents: After 48 h, the mice were euthanized by cervical dislocation.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Male C57BL/6J (6–8 weeks old, Harlan Laboratories, UK) were randomized into cages (n = 4/cage) and housed using 12 h dark/light cycles provided with water and fed ad libitum.
Subject Demographics
Age: Male C57BL/6J (6–8 weeks old, Harlan Laboratories, UK) were randomized into cages (n = 4/cage) and housed using 12 h dark/light cycles provided with water and fed ad libitum.
Randomization
Male C57BL/6J (6–8 weeks old, Harlan Laboratories, UK) were randomized into cages (n = 4/cage) and housed using 12 h dark/light cycles provided with water and fed ad libitum.
Blinding
not detected.
Power Analysis
Sample size calculations were based on previous variance (~15 %) between lean and obese mice in plasma 3H-cholesterol levels at 4h during RCT studies; $\mu1 = 0.458$; $\mu2 = 0.389$ and pooled variance $\sigma = 0.07$, where $N = 2\sigma2 (z\alpha/2 + z\beta)2/ (\mu1 – \mu2)2$.
Replication

not required.

Cell Line Authentication

Contamination: Cells were mycoplasma tested and IMPACT (IDEXX Laboratories, Inc., Maine, USA)-tested prior to injection.

Data Information

Availability: 2.7 Laboratory methods Detailed description of methodologies including glucose and insulin tolerance testing , lipoprotein analysis , acylcarnitine measurement , FPLC , real-time PCR , energy expenditure studies , stromal vascular fraction (SVF) isolation , and immunoblot analysis are available in the Supplemental Section .

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains			
Bodyweight and food consumption were measured weekly. 2.3 Rodent in vivo RCT studies Labeled J774.2 macrophages (European Collection of Cell Cultures) (~2 × 106 cells/4 × 106 dpm/mouse) were injected into C57BL/6J mice in the intraperitoneal cavity; the mice were housed individually in metabolic cages (Tecniplast, PA, USA) from 12 to 24 h post-injection and fed respective diets ad libitum.	C57BL/6J		Suggestion: RRID:IMSR_JAX:000664(link)
Software and Algorithms			
All other reagents, unless otherwise stated, were obtained from Sigma Aldrich Ltd. 2.2 Animals Animal care, procedures, and ethical approval were obtained from UCD, Ireland and UCC Ethics Committee.	Sigma Aldrich		Suggestion: Sigma-Aldrich (RRID:SCR_008988)(link)
Peptides and proteins were matched to the UniProt Human database and label-free quantitative (LFQ) ion intensities were generated using the default setting of MaxQuant , a quantitative proteomics software package.	UniProt		
	MaxQuant		Suggestion: MaxQuant (RRID:SCR_014485)(link)
Perseus statistical software was used to analyze the LFQ intensities .	Perseus		Suggestion: Perseus (RRID:SCR_015753)(link)
Differentially expressed proteins were inputted for pathway analysis through Ingenuity Pathway Analysis (IPA) and Cytoscape software.	Ingenuity Pathway Analysis		Suggestion: Ingenuity Pathway Analysis, RRID:SCR_008653(link)
	Cytoscape		Suggestion: Cytoscape (RRID:SCR_003032)(link)
GraphPad Prism version 5 (GraphPad Software Inc . , CA) was used for statistical analyses .	GraphPad Prism		
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
For comparison across multiple groups , data were tested for normal Gaussian distribution by the Shapiro–Wilk test .	Shapiro–Wilk test
Normally distributed raw or log-transformed data (n ≥ 8) , were tested by one-way or two-way analysis of variance (ANOVA) as appropriate and when significant Bonferroni post-hoc tests were applied.	analysis of variance
	ANOVA
	Bonferroni
Data-sets with n < 8 or non-normally distributed data underwent Kruskal–Wallis testing with Dunn 's post-hoc test .	Kruskal–Wallis testing
	Dunn 's post-hoc test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq